



### Organizing committee:

- ✓ Prof. Aleksandra Konjević, PhD, Prezident, Faculty of Agriculture, University of Novi Sad, Serbia
- ✓ Prof. Renata Bažok, PhD, Project Leader
- ✓ Prof. Mirjana Đukic-Stojčić, PhD, Vice dean for education and student activities, Faculty of Agriculture, University of Novi Sad, Serbia
- ✓ Prof. Bojan Konstantinović, PhD, Director of the Department for Environmental and Plant Protection, Faculty of Agriculture, University of Novi Sad, Serbia
- ✓ Goran Aleksić, PhD, President of the Plant Protection Society of Serbia
- ✓ Prof. Ferenc Bagi, PhD, Faculty of Agriculture, University of Novi Sad, Serbia
- ✓ Prof. Slavica Vuković, PhD, Faculty of Agriculture, University of Novi Sad, Serbia
- ✓ Prof. Sanja Lazić, PhD, Faculty of Agriculture, University of Novi Sad, Serbia
- ✓ Prof. Dušan Petrić, PhD, Faculty of Agriculture, University of Novi Sad, Serbia
- ✓ Prof. Vera Stojšin, PhD, Faculty of Agriculture, University of Novi Sad, Serbia
- ✓ Prof. Dragana Budakov, PhD, head of the Teaching chamber, Faculty of Agriculture, University of Novi Sad, Serbia
- ✓ Prof. Miloš Petrović, PhD, Faculty of Agriculture, University of Novi Sad, Serbia
- ✓ Doc. Mihaela Kavran, PhD, Faculty of Agriculture, University of Novi Sad, Serbia
- ✓ Marta Loc, PhD student, Faculty of Agriculture, University of Novi Sad, Serbia
- ✓ Milica Dudić, PhD student, Faculty of Agriculture, University of Novi Sad, Serbia
- ✓ Tatjana Dudaš, PhD student, Faculty of Agriculture, University of Novi Sad, Serbia
- ✓ Antonije Žunić, PhD student, Faculty of Agriculture, University of Novi Sad, Serbia
- ✓ Dragana Krajinović, Faculty of Agriculture, University of Novi Sad, Serbia
- ✓ Dejan Gligorić, Faculty of Agriculture, University of Novi Sad, Serbia
- ✓ Emil Šepecan, Faculty of Agriculture, University of Novi Sad, Serbia

### Supporters:

- ✓ Republic of Serbia, Autonomous Province of Vojvodina, Provincial Secretariat for Finance



## Content

**(According to Work Package 3 subgroups)**

## Subgroup 1 - Diagnosis in plant health and IPM

**Charikleia Kavroumatzi, Maria Iliadi, Maria Varveri, Anastasia Papageorgiou, Stavros Papageorgiou, Dimitrios Tsitsigiannis\*: Integrated management of *Alternaria dauci* in carrots..... 9**

**Ivana Miklečić<sup>1\*</sup>, Dana Čirjak<sup>1</sup>, Darija Lemić<sup>1</sup>, Tomislav Kos<sup>2</sup>, Ivana Pajač Živković<sup>1</sup>:**Detection of pear leaf blister moth using an automatic pest monitoring system ..... **10**

**Darija Lemic<sup>1</sup>, Mario Bjelis<sup>2</sup>, Helena Viric Gasparic<sup>1</sup>, Ivana Pajac Zivkovic<sup>1</sup>, Hugo A. Benitez<sup>3</sup>:**Geometric morphometrics techniques - a tool for evaluation of box tree moth *Cydalima perspectalis* invasiveness ..... **11**

**Maria Varveri<sup>1</sup>, Alex Bakos<sup>1</sup>, Panagiotis Gianniotis<sup>2</sup>, Dimitrios Tsitsigiannis<sup>1\*</sup>:**IKOPROTECTA  
- Agricultural composted products as plant protection and growth regulators..... **12**

## Subgroup 2 - Sustainable use of pesticides

**Nađa Milutinović<sup>1\*</sup>, Uroš Vojinović<sup>1</sup>, Tanja Vasić<sup>2</sup>, Milan Stević<sup>1</sup>:**Watermelon anthracnose disease in Serbia - importance and management..... **13**

**Helena Viric Gasparic<sup>1\*</sup>, Sanja Lazić<sup>2</sup>, Aleksandra Šušnjar<sup>2</sup>, Dragana Šunjka<sup>2</sup>:**Pesticide residue analysis of plant samples..... **14**

**Dana Čirjak<sup>1\*</sup>, Ivana Miklečić<sup>1</sup>, Darija Lemić<sup>1</sup>, Tomislav Kos<sup>2</sup>, Ivana Pajač Živković<sup>1</sup>:** Overview of automatic monitoring systems for apple pests ..... 15

**Evgenia-Anna Papadopoulou<sup>1</sup>, Alexios-Leandros Skaltsounis<sup>2</sup>, Konstantinos A. Aliferis<sup>1,3\*</sup>:**Olive tree *endophytes*: a new source of bioactivity in plant protection ..... 16

**Uroš Vojinović\*, Milan Stević:**Current status of *Erysiphe necator* populations sensitivity to fungicides in Serbia ..... 17

**Dragan Jurković<sup>1</sup>, Mladen Zovko<sup>1</sup>, Vjekoslav Tadić<sup>2</sup>:**Influence of different spraying technical factors on leaf surface coverage in vineyard ..... **18**

**Milica Dudić<sup>1\*</sup>, Maja Meseldžija<sup>1</sup>, Verica Vasić<sup>2</sup>:**Systemic herbicides use in poplar plantations ..... 19

Bajagić, B., Latinović, J. and N. Latinović: Significance of Appropriate Pesticide Application 20

**Marko Sretenović\*, Novica Miletić, Nenad Tamaš:**Efficacy of different insecticides in the

control of green peach aphid on nectarine..... 21

**Sandra Skendžić<sup>1</sup>\*, Monika Zovko<sup>1</sup>, Darija Lemić<sup>1</sup>, Nadja Štetin<sup>2</sup>, Branislav Jović<sup>2</sup>:**Soil reflectance spectroscopy - a tool for determining and monitoring soil properties ..... 22

### Subgroup 3 - Plant feeders

**Renata Bažok<sup>1</sup>\*, Elizabeta Gavić<sup>1</sup>, Tena Krajinović<sup>1</sup>, Siniša Jelovčan<sup>2</sup>, Maja Čačija<sup>1</sup>:**Effects of spiropidion on survival and parasitism of *Aphidius collemanni* Viereck (*Hymenoptera: Braconidae*) ..... 23

**Martina Kadoić Balaško<sup>1</sup>, Sandra Senji<sup>2</sup>, Aleksandra Konjević<sup>2</sup>:**Electrical penetration graph (EPG) – promising tool for measuring feeding behaviour of *Bemisia tabaci*..... 24

**Ilaria Laterza\*, Enrico de Lillo, Giovanni Tamburini:**Impacts of landscape composition on the occurrence of *Halyomorpha halys* and biological control potential in newly invaded areas.. 25

**Mariya Hristozova:**Parasitoids of *Halyomorpha halys* (Heteroptera: Pentatomidae) in Bulgaria..... 26

**Sofia A. Dervisoglou<sup>\*1</sup>, Dionysios Ch. Perdakis<sup>1</sup>, Nikos E. Papanikolaou<sup>1,2,†</sup>, Argyro A. Fantinou<sup>3</sup>:***Tuta absoluta* egg density and distribution on tomato plants affect the efficacy of mirid predators..... 27

**Ivana Lalićević\*, Anđa Radonjić:**Aphids flight activity on wheat..... 28

**Marija Milošević<sup>1</sup>\*, Aleksandar Stojanović<sup>2</sup>, Draga Graora<sup>1</sup>:**Natural enemies of *Parthenolecanium corni* (Bouché) (Hemiptera: Coccidae) on grapevine in Neštin, Serbia ..... 29

**Mihaela Kavran, Aleksandra Konjević, Aleksandra Ignjatović Čupina, Miloš Petrović, Dušan Petrić:**Some important invasive species in agricultural and urban environments of Serbia ..... Error! Bookmark not defined.

**Gianvito Ragone:**Sustainable control of insect pests in Mediterranean agroecosystems..... 31

**Aleksandar Ivezić\*, Aleksandra Ignjatović Čupina:**Identification of *Trichogramma* egg parasitoids of the European corn borer *Ostrinia nubilalis* (Hübner, 1796) in Vojvodina (Serbia) ..... 32

**Vjekoslav Markotić<sup>1</sup>\*, Renata Bažok<sup>2</sup>, Tatjana Masten Milek<sup>3</sup>, Mladen Šimala<sup>1</sup>:** Fauna of insects from the suborder Sternorrhyncha in citrus orchards in Dubrovačko -neretvanska County.....33

### Subgroup 4 - Plant Pathology

**Rita Milvia De Miccolis Angelini<sup>1</sup>, Celeste Raguseo<sup>1</sup>, Lucia Landi<sup>2</sup>, Caterina Rotolo<sup>1</sup>, Palma Rosa Rotondo<sup>1</sup>, Donato Gerin<sup>1</sup>, Gianfranco Romanazzi<sup>2</sup>, Stefania Pollastro<sup>1</sup>, Francesco**

- Faretra<sup>1\*</sup>**:Multi-omics approaches to explore phenotypic and genetic diversity in the brown rot fungal pathogens *Monilinia fructicola*, *Monilinia laxa* and *Monilinia fructigena* ..... 34
- Tatjana Dudaš<sup>1\*</sup>, Dragana Budakov<sup>1</sup>, Simona Sanzani<sup>2</sup>, Mila Grahovac<sup>1</sup>, Marta Loc<sup>1</sup>, Vera Stojšin<sup>1</sup>, Antonio Ippolito<sup>2</sup>**:Biocontrol activity of epiphytic yeasts against *Penicillium expansum* on apple fruits..... 35
- Konjić Almira\*, Arnela Okić, Jasmin Grahić, Fuad Gaši**:Overview of the use of GWAS to identify loci for resistance to apple scab caused by *Venturia Inaequalis* ..... 37
- Anastasia Papageorgiou, Maria Varveri, Dimitrios Tsitsigiannis\***:Endophytic bacteria from olive drupes as biological control and induced resistance agents of *Colletotrichum acutatum* ..... 38
- Eirini Poulaki<sup>1\*</sup>, Ioannis Stringlis<sup>2</sup>, Marinos Xagoraris<sup>3</sup>, Christos Pappas<sup>3</sup>, Sotiris Tjamos<sup>1\*</sup>**:The volatile organic compounds of the biocontrol agent *Bacillus velezensis* K165 activate plant defenses against *Verticillium dahliae*..... 39
- Christos Tsoukas, Aliko Tzima, Epaminondas Paplomatas<sup>1</sup>**:Grapevine rootstock and scion canes as an important inoculum source of Petri disease associated pathogens ..... 40
- Nina Vučković\*, Ivana Vico, Nataša Duduk**:Botryosphaeriaceae as postharvest pathogens of apple fruit ..... 41
- Polina Tsalgatidou<sup>1,2</sup>, Eirini Evangelia Thomludi<sup>1</sup>, Eirini Baira <sup>3</sup>, Panagiotis Katinakis<sup>1</sup>, Anastasia Venieraki<sup>4\*</sup>**:Antagonistic activity of endophytic *Bacillus* strains against phytopathogenic fungi. Mode of action studies and the strain compatibility issue ..... 42
- Saud Hamidović\*, Fejzo Bašić, Sanel Haseljić, Teofil Gavrić, Arnela Okić, Berina Imamović, Nedžad Karić**:Antifungal activity of peppermint, sweet basil, and sage essential oils on plant pathogenic fungi..... 43
- Magdalena Matic\***:Physiological response in winter wheat varieties depending on *Fusarium* infection and N fertilization ..... 44
- M. Mastrochirico, Roberta Spanò, Tiziana Mascia\***:Grafting as a sustainable approach in response to infections transmitted by arthropods..... 45
- Palma Rosa Rotondo<sup>1</sup>, Rita Milvia De Miccolis Angelini<sup>1\*</sup>, Paolo Francesco Ambrico<sup>2</sup>, Caterina Rotolo<sup>1</sup>, Donato Gerin<sup>1</sup>, Stefania Pollastro<sup>1</sup>, Francesco Faretra<sup>1</sup>**:Cold Plasma as a new technology in the sustainable management of plant diseases..... 46
- Charikleia Kavroumatzi, Maria Iliadi, D. Akrivopoulou, Dimitrios Tsitsigiannis\***:Evaluation of biological and synthetic plant protection products for the management of downy mildew in





production in *Penicillium* spp. causing postharvest pome fruit decay .....587

**Tamara Siber\***:Antifungal action of quaternary pyridinium compounds on phytopathogenic polyphagous fungi species of the genera *Fusarium*, *Botryotinia* and *Sclerotinia*.....598

**Ana Sesar<sup>1</sup>, Anita Lalić<sup>2\*</sup>, Kristina Batinić<sup>1</sup>**:Occurrence of ochratoxin A in Blatina wine from the Mostar vineyards .....609

**Maria Varveri, Mike Kaminiaris, Spyridoula Dimitropoulou, Dimitrios Tsitsigiannis\***:  
Biological control of aflatoxins using non-toxigenic strains of *Aspergillus flavus* on maize.... 60

### Subgroup 7 - Bio-diversity and bio-indicators in sustainable agriculture

**Robert Damo, Pirro Icka\***:Neglected and Underutilized Crop Species, an alternative to improve food supply and nutrition security by protecting biodiversity in Korça region, Albania.....621

**Josipa Puškarić<sup>1\*</sup>, Vladimir Ivezić<sup>1</sup>, Brigita Popović<sup>1</sup>, Dinka Grubišić<sup>2</sup>, Milan Radivojević<sup>3</sup>, Nikola Grujić<sup>3</sup>, Mirjana Brmež<sup>1</sup>**:Nematodes as bioindicators of soil ecosystem status in the intercropping system of woody species and agricultural crops .....632

**Program of the conference**.....63





## Subgroup 1 - Diagnosis in plant health and IPM

### Detection of pear leaf blister moth using an automatic pest monitoring system

Ivana Miklečić<sup>1\*</sup>, Dana Čirjak<sup>1</sup>, Darija Lemić<sup>1</sup>, Tomislav Kos<sup>2</sup>, Ivana Pajač Živković<sup>1</sup>

<sup>1</sup> University of Zagreb Faculty of Agriculture, Department of Agricultural Zoology,  
Svetosimunska cesta 25, 10 000 Zagreb, Croatia

<sup>2</sup> University of Zadar, Department of Ecology, Agriculture and Aquaculture, Trg kneza  
Višeslava 9, 23 000 Zadar, Croatia

\*Corresponding author: [imiklecic@agr.hr](mailto:imiklecic@agr.hr)

**Abstract:** Pear leaf blister moth (Lepidoptera: Lyonetiidae) is an economically dangerous insect in apple orchards worldwide. During the growing season, the moth develops several generations that cause damage directly to the leaves. The larvae live inside the leaves and feed on the mesophyll tissue, causing defoliation of the leaves and later affecting bud differentiation. In more severe infestations, fruit organoleptic characteristics may also be affected. In the context of precision agriculture, the introduction of new technologies and early detection of pests through the use of automated monitoring systems for economically important apple pests can improve pest control and reduce damage to apple crops. The objective of this study is to develop an automatic monitoring system based on pear leaf blister moth detection using an RGB camera. During a period of 11 weeks, a series of 250 images were taken, and 4150 moths were annotated in the images. A convolutional neural network was trained based on 90% of the annotated images. The results showed that the analytical model trained in this way was able to identify the pear leaf blister moth with 70% accuracy, which was verified using the remaining 10% of the captured images. These preliminary results show that the system can contribute to more accurate and early detection of pests, but still needs to be improved with a larger training data set.

**Keywords:** *Leucoptera malifoliella* (Costa 1836), apple production, artificial neural network, precision agriculture

## Subgroup 1 - Diagnosis in plant health and IPM

### Geometric morphometrics techniques - a tool for evaluation of box tree moth *Cydalima perspectalis* invasiveness

Darija Lemic<sup>1</sup>, Mario Bjelis<sup>2</sup>, Helena Viric Gasparic<sup>1</sup>, Ivana Pajac Zivkovic<sup>1</sup>, Hugo A. Benitez<sup>3</sup>

<sup>1</sup> University of Zagreb Faculty of Agriculture, Department of Agricultural Zoology,  
Svetosimunska 25, 10000 Zagreb, Croatia ([hvirc@agr.hr](mailto:hviric@agr.hr))

<sup>2</sup> University of Split, University Department of Marine Studies, Ruđera Boskovića 37, 21000,  
Split, Croatia

<sup>3</sup> Laboratorio de Ecología y Morfometría Evolutiva, Centro de Investigación de Estudios  
Avanzados del Maule, Universidad Católica del Maule, Talca, Chile

\*Corresponding author: [dlemic@agr.hr](mailto:dlemic@agr.hr)

**Abstract:** Morphological variation can affect the ability of species to disperse and adapt to new areas. Geometric morphometric methods are valuable tools for assessing the extent of phenotypic changes influenced by genetic variation. Such methods were used in this study to determine the morphological variability of the box tree moth *Cydalima perspectalis*, Walker from Croatia and to assess its invasive character. Box tree moth is an invasive species that spreads rapidly, has high reproductive potential, and is highly adaptable. A cross-site analysis was conducted, and the dataset was partitioned by geographic area, with intraspecific variation found independently between northern and southern populations. Results indicate that wing shape is influenced by agroecological factors, with forewings being more plastic than hindwings. Northern populations have broader wings, while southern populations have more elongate wings. Wing characteristics of box tree moth are of great importance in assessing their spread, especially given their high invasive potential. Using this innovative and effective, yet simple and inexpensive method, we can detect differences in wing shape and size and use these differences as biomarkers of variation between and within populations. These results suggest that geometric morphometric methods can be used to study the biological adaptation of box tree moth in native and newly invaded areas.

**Keywords:** box tree moth, landmarks, adaptation, wings, geometric morphometrics, invasiveness

## Subgroup 1 - Diagnosis in plant health and IPM

### IKOPROTECTA - Agricultural composted products as plant protection and growth regulators

Maria Varveri<sup>1</sup>, Alex Bakos<sup>1</sup>, Panagiotis Gianniotis<sup>2</sup>, Dimitrios Tsitsigiannis<sup>1\*</sup>

<sup>1</sup> Laboratory of Plant Pathology, Department of Crop Science

Agricultural University of Athens, Greece, <sup>2</sup>IKORGANIC, Kifissia, Greece

\*Corresponding author: [dimtsi@aia.gr](mailto:dimtsi@aia.gr)

**Abstract:** Composts are natural products known for their beneficial properties in plant nutrition by enriching the soil with organic matter and nutrients. Composts contain high microbe populations, that are mainly non-pathogenic and a number of studies have highlighted their capacity to suppress or control a variety of plant diseases. The aim of this study was to assess the biopotential of two composted products (Ikorganic), based on disease assessments in greenhouse and *in vitro* experiments. The fungi, yeasts and bacteria that were isolated from the composted products, were evaluated on their ability to inhibit the growth of phytopathogenic fungi and bacteria such as *Verticillium dahliae*, *Botrytis cinerea* and *Pseudomonas syringae* pv tomato. The antagonistic activity of the isolates was investigated, *in vitro*, by employing dual culture technique and several of efficient antagonistic bacterial isolates were identified. Subsequently, the efficiency of the composted products concerning the plant growth promotion and the reduction of disease severity against the above-mentioned pathogens, was evaluated and demonstrated in greenhouse, on tomato plants and on tomato fruits and strawberries. The results of the *in vitro* experiments as well as the comparative study of the two different composted products regarding their ability to control both foliar and soil-borne plant diseases will be presented.

**Keywords:** composts, plant protection, biological control, plant growth promotion

*Acknowledgment:* This research has been co-financed by the European Regional Development Fund of the European Union and Greek national funds through the Operational Program Competitiveness, Entrepreneurship and Innovation, under the call RESEARCH – CREATE – INNOVATE (project code: T2EΔK-00965, IKOPROTECTA)

## Subgroup 2 - Sustainable use of pesticides

### Watermelon anthracnose disease in Serbia - importance and management

**Nađa Milutinović<sup>1\*</sup>, Uroš Vojinović<sup>1</sup>, Tanja Vasić<sup>2</sup>, Milan Stević<sup>1</sup>**

<sup>1</sup>University of Belgrade, Faculty of Agriculture, Belgrade-Zemun, Serbia

<sup>2</sup>University of Niš - Faculty of Agriculture, Kruševac, Serbia

\* Corresponding author: [nadiaamilutinovic10@gmail.com](mailto:nadiaamilutinovic10@gmail.com)

**Abstract:** Watermelon (*Citrullus lanatus* (Thunb.) Matsum. et Nakai) is one of the most important cucurbit crop worldwide. Watermelon is generally grown in open-fields, where the environmental conditions are favorable for disease development. Anthracnose, caused by the fungal pathogen *Colletotrichum orbiculare* (Berk. et Mont.) Arx (syn. *C. lagenarium* (Pass.) Ellis et Halst.), is a destructive foliage disease in watermelon production. In warm, humid climates around the world, the fungus affects watermelon and many other cucurbits, and it can lead to severe crop losses. The disease significantly reduces the total amount of yield and fruit quality by infecting above-ground organs like fruits, leaves, and stems. Due to that reason, effective control of *C. orbiculare* has to be done. It can be achieved using all preventive measures, whereas the use of fungicides has a major importance. However, management of anthracnose is very difficult because there are no registered fungicides in the Serbian pesticide market for chemical control of *C. orbiculare*. Considering that the watermelon is a “minor crop”, the registration of plant protection products is not profitable for agrochemical industry. Based on the results of morphological, pathogenic and molecular characterization, all isolates used in this study were determined as *C. orbiculare*. Using the *in vitro* and *in vivo* methods, the sensitivity of *C. orbiculare* isolates to different fungicides were established in laboratory.

**Keywords:** *Colletotrichum orbiculare*, watermelon, anthracnose, fungicides, sensitivity.

## Subgroup 2 - Sustainable use of pesticides

### Pesticide residue analysis of plant samples

Helena Viric Gasparic<sup>1\*</sup>, Sanja Lazić<sup>2</sup>, Aleksandra Šušnjar<sup>2</sup>, Dragana Šunjka<sup>2</sup>

<sup>1</sup>University of Zagreb Faculty of Agriculture, Svetošimunska 25, Zagreb, Croatia

<sup>2</sup>University of Novi Sad Faculty of Agriculture, Trg Dositeja Obradovica, 8, Novi Sad, Serbia

\* Corresponding author: [hvirc@agr.hr](mailto:hviric@agr.hr)

**Abstract:** Plant protection products are widely used in agricultural production to control pests, increase yield, improve quality, and extend the shelf life of produce. Although their properties are usually valuable, these very properties can have negative effects when pesticide residues are released into the environment. The presence of pesticide residues, their metabolites or degradation products in the environment is a significant risk factor. Therefore, out of concern for public health, maximum residue limits (MRLs) were established by Regulation (EC) No. 396/2005. MRLs are legally permissible upper limits for pesticide residues in or on food or animal feed based on good agricultural practice (GAP) and the lowest exposure necessary to protect vulnerable consumers. Residue analysis provides a measure of the nature and extent of chemical contamination in the environment and its persistence. With the increasing focus on food safety, accurately determining pesticide residues in food has become an important issue. The most common analytical techniques for the detection of pesticide residues are gas chromatography-mass spectrometry (GC-MS) for volatile compounds and liquid chromatography-mass spectrometry (LC-MS) for non-volatile, thermally unstable, and polar compounds without derivative. The combination of chromatography and MS is become one of the most powerful tools for screening, confirmation, and quantification of complicated organic contaminants. The authors will present the advantages of each method as an important procedure for determining the safety of using certain pesticides.

**Keywords:** analytical chemistry, detection, food safety, pesticides, residues.

## Subgroup 2 - Sustainable use of pesticides

### Overview of automatic monitoring systems for apple pests

Dana Čirjak<sup>1\*</sup>, Ivana Miklečić<sup>1</sup>, Darija Lemić<sup>1</sup>, Tomislav Kos<sup>2</sup>, Ivana Pajač Živković<sup>1</sup>

<sup>1</sup> University of Zagreb Faculty of Agriculture, Department of Agricultural Zoology,  
Svetošimunska cesta 25, 10 000 Zagreb, Croatia

<sup>2</sup> University of Zadar, Department of Ecology, Agriculture and Aquaculture, Trg kneza  
Višeslava 9, 23 000 Zadar, Croatia

\*Corresponding author: [dcirjak@agr.hr](mailto:dcirjak@agr.hr)

**Abstract:** Apple (*Malus domestica* Borkh.) is one of the world's highest-yielding fruit crops. In addition to codling moth (*Cydia pomonella* L.), which is considered the most important apple pest, other lepidopterans, fruit flies, and true bug species also threaten production, affecting fruit quality and preventing its commercialization. Early detection and monitoring of pests is critical to preventing damage from pests. Due to new climatic conditions, the phenology of pests changes frequently and it is very difficult to predict their occurrence. Therefore, there is an increasing need to develop and use new smart technologies for pest monitoring (smart traps, sensors, etc.). Such methods can greatly facilitate site-specific management and consequently reduce the use of pesticides and their residues. The aim of this work is to provide an overview of methods for automatic monitoring of economically important apple pests in order to improve sustainable pest management in apple production.

**Keywords:** apple production, smart traps, site-specific management



## Subgroup 2 - Sustainable use of pesticides

### Olive tree endophytes: a new source of bioactivity in plant protection

Evgenia-Anna Papadopoulou<sup>1</sup>, Alexios-Leandros Skaltsounis<sup>2</sup>, Konstantinos A. Aliferis<sup>1,3\*</sup>

<sup>1</sup>Agricultural University of Athens, Laboratory of Pesticide Science, Iera Odos 75, Athens, 118 55

<sup>2</sup>Department of Pharmacognosy and Natural Products Chemistry, Faculty of Pharmacy, University of Athens, Athens, Greece

<sup>3</sup>McGill University, Department of Plant Science, 21111 Lakeshore Road, H9X 3V9, Montreal, QC, Canada

\*Corresponding author: [konstantinos.aliferis@aua.gr](mailto:konstantinos.aliferis@aua.gr); [evina.papadopoulou@gmail.com](mailto:evina.papadopoulou@gmail.com)

**Abstract:** The agri-food sector is facing unprecedented challenges, which necessitate innovative approaches and methods in order to be addressed. Among them, climate change, the emerging food crisis due to the recent war, and plant protection issues threaten the supply of the world's population with sufficient quantities of food. However, the application of traditional farming models and research based on conventional methodologies and approaches are not sufficient to solve the aforementioned issues. Therefore, here, a new source of bioactivity for applications in plant protection, olive tree endophytes, was assessed. Endophytes can be used in plant protection *per se*, or their bioactive metabolites could be potentially developed as pesticides. In this context, a holistic approach was applied for the isolation of olive tree endophytes and the assessment of their bioactivity. The developed pipeline resulted in the isolation of a large number of fungal and bacterial endophytes. Following the bioactivity assessment, a bacterial isolate exhibited superior antifungal activity against the devastating *Colletotrichum acutatum* species complex. Such bioactivity was largely attributed to the production of lipopeptides, whose toxicity was further assessed to target and non-target organisms and their plant-priming capacity was assessed *in planta* to olive tree applying metabolomics. The results were promising towards the further development of the bacterium as a plant protection agent, and the established pipeline could be used in the R&D of endophytic microorganism-based pesticides, that could, at least partially, substitute conventional pesticides.

**Keywords:** fungicides, metabolomics, natural products

## Subgroup 2 - Sustainable use of pesticides

### Current status of *Erysiphe necator* populations sensitivity to fungicides in Serbia

Uroš Vojinović\*, Milan Stević

Faculty of Agriculture – University of Belgrade, Nemanjina 6, Zemun-Belgrade, Serbia

\*Corresponding author: [uros.vojinovic@agrif.bg.ac.rs](mailto:uros.vojinovic@agrif.bg.ac.rs)

**Abstract:** The grapevine production in Serbia takes place on approximately 21000 hectares, with an annual total production of about 160000 tons. One of the most important pathogen in grapevine cultivation is *Erysiphe necator*, the causal agent of powdery mildew. If chemical protection is not carried out properly, the pathogen may cause significant losses in the yield and quality of the grapes. Systemic fungicides are prevalently used to control *E. necator*. Due to their very specific mode of action and continuous application, the risk of resistance development is increased. The results about this topic in Serbia are scarce, so the aim of this study was to investigate the efficacy of different fungicides and determine the sensitivity status of *E. necator* populations. The trials were conducted on four locations from 2019 to 2021. Completely randomized block system with four replications was used. Commercial formulations of the fungicides from different groups (QoI, SDHI, DMI fungicides, spiroketal-amines, aryl-phenyl ketones, aza-naphtalenes, phenyl-acetamide) were tested. During each experimental year, two assessments were done on the bunches and one on the leaves. The incidence and severity of powdery mildew in the control plots were significant especially during 2019 and 2020. A drop in the efficacy of QoI fungicides and DMI fungicides was observed. Evidently reduced efficacy of aryl-phenyl-ketones was observed on two localities. A lower sensitivity of the population to quinoxifen was determined, but no cross resistance between aza-naphtalenes was detected. In two experimental sites, the efficacy of boscalid was low, while the other SDHI fungicides showed very high efficacy in all experimental sites. Depending on the site, the efficacy of spiroxamine and cyflufenamide was moderately high to very high. The results obtained in this study indicate that there has been a shift in the sensitivity of *E. necator* populations to certain groups of fungicides in Serbia.

**Keywords:** powdery mildew, grapevine, sensitivity, fungicides, efficacy



## Subgroup 2 - Sustainable use of pesticides

### Systemic herbicides use in poplar plantations

Milica Dudić<sup>1\*</sup>, Maja Meseldžija<sup>1</sup>, Verica Vasić<sup>2</sup>

<sup>1</sup>Faculty of Agriculture, University of Novi Sad, Dositej Obradović Square 8, 21000 Novi Sad, Serbia

<sup>2</sup>Institute of Lowland Forestry and Environment, University of Novi Sad, Antona Čehova 13d, 21000 Novi Sad, Serbia

\* Corresponding author: [milica.dudic@polj.edu.rs](mailto:milica.dudic@polj.edu.rs)

**Abstract:** The aim of this study was to determine the presence of weeds and to investigate the efficacy of the herbicides glyphosate (6 l/ha) and triclopyr (4 l/ha) in three-year-old poplar plantations. A field study was conducted in 2018 and 2019, at the Institute of Lowland Forestry and Environment “Kačka forest”. The experiment was established as a random block design with four replications. Herbicide efficacy and phytotoxicity were evaluated after 15 and 30 days. In 2018, the presence of 19 weed species was identified (*Amorpha fruticosa*, *Apera spica-venti*, *Apium graveolens*, *Asclepias syriaca*, *Avena fatua*, *Cirsium arvense*, *Convolvulus arvensis*, *Dactylis glomerata*, *Equisetum arvense*, *Galium lanceolatum*, *Geranium dissectum*, *Glechoma hederacea*, *Ranunculus repens*, *Solidago gigantea*, *Solidago speciosa*, *Symphytum officinale*, *Trifolium campestre*, *Trifolium repens*, *Vicia villosa*), of which 84,21% belong to broad-leaved species and 15,79% to grass species. During 2019, the presence of 18 weed species was identified (*Amorpha fruticosa*, *Apera spica-venti*, *Asclepias syriaca*, *Avena fatua*, *Capsella bursa-pastoris*, *Dactylis glomerata*, *Daucus carota*, *Erigeron annuus*, *Equisetum arvense*, *Galium lanceolatum*, *Glechoma hederacea*, *Solidago gigantea*, *Solidago speciosa*, *Symphytum officinale*, *Trifolium campestre*, *Trifolium repens*, *Veronica hederifolia* and *Vicia villosa*), of which 83.33% belong to broad-leaved species and 16.67% to grass weeds. Results show that glyphosate had the best efficacy which was 94.34%, while the total efficacy of triclopyr was 85.53%. Herbicide phytotoxicity was not recorded in three-year poplar plantations. The high potency of controlling weeds showed that glyphosate and triclopyr can be recommended as an effective measure to control weeds in poplar plantations.

**Keywords:** weed vegetation, glyphosate, triclopyr, weed control, poplar

## Subgroup 2 - Sustainable use of pesticides

### Significance of Appropriate Pesticide Application

**Bajagić, B., Latinović, J. and N. Latinović**

*University of Montenegro, Biotechnical Faculty, Mihaila Lalica 15, 81000, Podgorica,  
Montenegro*

\* Corresponding author: [balsab@ucg.ac.me](mailto:balsab@ucg.ac.me)

**Abstract:** In order to successfully protect cultivated plants from harmful organisms, it's necessary not only to choose an appropriate plant protection product and to use it in a timely manner, but also to properly apply it to the plant. Therefore, the appropriate choice of equipment for application, it's proper functioning have a significant impact on efficiency of the treatment. In accordance with the Directive of the European Parliament 2009/128 EC, frameworks on the sustainable use of plant protection products have been established. Inspection of pesticide application equipment includes checking the technical accuracy and functionality of the device as well as risk assessment for the operator and the environment. The standardization of this procedure in Europe was established in 2003 with the adoption of the European standard EN 13790, and by 2006 it's implementation became mandatory in 20 countries.

In Montenegro, during 2021, the Directorate for Food Safety, Veterinary and Phytosanitary Affairs implemented a national program of phytosanitary measures, which prescribed the inspection of equipment used for the application of plant protection products. The objective was to get an overview on condition of the equipment for pesticide application in Montenegro. Major irregularities were observed with most of the devices, such as malfunctioning of manometers and uneven flow of nozzles, which directly affects the uncontrolled application of pesticides. After testing of 20 devices for pesticide application, only one met the technical correctness. In further activities, it's planned to intensify the control of pesticide application equipment in order to achieve controlled use of pesticides.

**Keywords:** equipment, sustainable use of pesticides

## Subgroup 2 - Sustainable use of pesticides

### Efficacy of different insecticides in the control of green peach aphid on nectarine

Marko Sretenović\*, Novica Miletić, Nenad Tamaš

University of Belgrade, Faculty of Agriculture, Zemun

\* Corresponding author: [sretenovic@agrif.bg.ac.rs](mailto:sretenovic@agrif.bg.ac.rs)

**Abstract:** The efficacy of different insecticides in the control of green peach aphid (*Myzus persicae*) on nectarine at the the locality of Metković (Bogatić municipality) was examined conducting field trials in 2019 and 2020. The experiments was carried out according to the statistical model of completely randomized block design with four replications and standard EPPO method for evaluation of the efficacy of insecticides in the control of aphids in orchards. Insecticidal formulations were previously diluted with water and applied with a backpack sprayer "Solo". The efficacy of insecticides from the group of neonicotinoids (acetamiprid, imidacloprid, thiamethoxam), organophosphates (chlorpyrifos), pyrethroids (deltamethrin) and flonicamid were examined. Results of the experiments indicate very good efficacy of insecticides from the group of neonicotinoids in the control of *M. persicae*. The efficacy of acetamiprid three days after treatment (3DAT) was 98.23%, while seven days after treatment (7DAT) it was 99.43%. Imidacloprid showed maximum efficacy (100%) in both evaluation terms (3DAT and 7DAT). The efficacy of thiamethoxam was high, 99.69% (3DAT) and 99.92% (7DAT), respectively. Deltamethrin showed efficacy of 61.37% (3DAT) and its efficacy was 47.17% at 7DAT. The efficacy of chlorpyrifos ranged from 31.75% (7DAT) to 68.84% (3DAT). Flonicamid, considering the mode of action (anti-feeding), had a lower initial efficacy (82.13%), while the efficacy at 7DAT was significant higher (99.65%).

**Keywords:** *M. persicae*, nectarine, field trials, insecticides, efficacy



## Subgroup 2 - Sustainable use of pesticides

### Soil reflectance spectroscopy - a tool for determining and monitoring soil properties

Sandra Skendžić<sup>1</sup> \*, Monika Zovko<sup>1</sup> , Darija Lemić<sup>1</sup> , Nadja Štetin<sup>2</sup> , Branislav Jović<sup>2</sup>

*1 University of Zagreb Faculty of Agriculture, Svetosimunska 25, 10000 Zagreb, Croatia*

*2 University of Novi Sad, Faculty of Sciences, Dositeja Obradovica 3, Novi Sad 21000, Serbia*

\* Corresponding author: [sskendzic@agr.hr](mailto:sskendzic@agr.hr)

**Abstract:** Intensive agriculture has slowly degraded soils through pollution with pesticide residues, nutrients, and salinization, as well as crop residue removal, loss of soil organic carbon, and other threats. Therefore, a loss of soil fertility is observed. Soil analysis is an essential tool for precise management of fertilisation and irrigation, which enables to obtain high quality crops. Soil spectroscopy has proven to be a rapid, cost-effective, environmentally friendly, non-destructive, and analytical technique for determining soil properties. The main objective of this study was to propose an effective approach based on FTIR spectroscopy for determining soil properties of soil samples from different sites and management regimes. Spectral signatures in the mid-infrared range (4000-700 cm<sup>-1</sup>) were acquired using a spectrophotometer, and chemometric methods were used to correlate chemical and physical soil properties with their spectral signatures. Various calibration models can be used to quantitatively predict soil properties - such models were developed as a function of measured data from soil laboratory analyses and soil reflectance spectra. The FTIR spectra of the soil samples were subjected to multivariate analysis in the form of principal component analysis (PCA) and cluster analysis (CA). The multivariate analysis of the FTIR soil spectral data showed that it is possible to distinguish and classify soils at different sites based on their spectral signatures. These results demonstrate that FTIR spectroscopy and multivariate analysis can be used as a tool to identify and classify soil samples.

**Keywords:** soil properties, FTIR spectroscopy, reflectance, agriculture



### Subgroup 3 - Plant feeders

## Electrical penetration graph (EPG) – promising tool for measuring feeding behaviour of *Bemisia tabaci*

Martina Kadoić Balaško<sup>1</sup>, Sandra Senji<sup>2</sup>, Aleksandra Konjević<sup>2</sup>

<sup>1</sup>University of Zagreb, Faculty of Agriculture, Svetosimunska 25, Zagreb, Croatia

<sup>2</sup>University of Novi Sad, Faculty of Agriculture, Trg Dositeja Obradovica, 8, Novi Sad, Serbia

\* Corresponding author: [mbalasko@agr.hr](mailto:mbalasko@agr.hr)

**Abstract:** *Bemisia tabaci* (Sweet potato whitefly) is one of the most important pests in greenhouse production systems worldwide. It causes great damage to vegetables at all stages of development. It is highly adaptable to new hosts and climatic conditions, which is why it has been found on 300 different hosts and reported from all over the world. *B. tabaci* belongs to the herbivorous hemipteran insects that penetrate plant tissue with their piercing-sucking mouthparts. They damage plants by depriving them of photosynthetic substances as they feed, by excreting large amounts of honeydew on which saprophytic fungi spoil leaves and fruits, and by transmitting yield-reducing viruses. Therefore, it is important to know the feeding habits and preferences of this pest. A useful tool for studying the feeding habits of sucking insects is the electrical penetration graph (EPG). In an EPG measurement, an electrical circuit is established between the plant and the insect when the insect contact the plant. Thus, using this method, it is possible to visualize the feeding behavior of hemipteran's in real time. A good knowledge of insect-plant interactions is necessary to identify and evaluate resistant traits for selection in plant breeding programs. In our research we used tomato leaves, which were treated with clay and water to see the differences in feeding habits. We will present preliminary results of the EPG as a promising tool to measure the feeding behavior of *B. tabaci*. Our preliminary data will be used in future research on the feeding behavior of this important pest.

**Keywords:** insect-plant interaction, sucking insects, feeding behavior, insect pest, *Bemisia tabaci*

### Subgroup 3 - Plant feeders

## Impacts of landscape composition on the occurrence of *Halyomorpha halys* and biological control potential in newly invaded areas

Ilaria Laterza\*, Enrico de Lillo, Giovanni Tamburini

University of Bari Aldo Moro - Department of Soil, Plant and Food Sciences (DiSSPA).

\* Corresponding author: [ilaria.laterza@uniba.it](mailto:ilaria.laterza@uniba.it)

**Abstract:** Early understanding of spreading pathways of alien pests in newly invaded areas is crucial to plan efficient strategies to prevent serious outbreaks. *Halyomorpha halys* is an East Asian invasive species, now established across Europe and North America, causing important economic losses. The pest was recorded for the first time in Northern Italy in 2012 and in 2016 in the Apulia region (Southern Italy), spreading across the Italian Peninsula, often as hitchhiker following commercial trade ways. Information regarding its occurrence and factors shaping its population dynamics in Southern Italy remain scarce. In this study we investigated the occurrence of *H. halys* in orchards, urban gardens, and fruit warehouses habitat in the Apulia region. Sites were selected along a gradient of urbanization in the landscape. We also measured the control potential provided by parasitoids and generalist predators exposing refrigerated egg masses in orchards and urban gardens. Our results showed no differences in the occurrence and abundance of the pest across habitats. We found a positive effect of urban area in the landscape on the occurrence and abundance of the pest in orchard habitat. We also observed significant positive effect of seminatural habitat on the percentage of parasitized and preyed eggs. These preliminary results suggest that the *H. halys* can use multiple habitats and that its spreading is linked to the presence of human settlements, while seminatural habitats play a key role in supporting the biological control. Our findings might be useful to plan more efficient monitoring and control strategies able to limit its further spread.

**Keywords:** Brown Marmored Stink Bug, invasive pest, monitoring program, pest dynamics, landscape composition

### Subgroup 3 - Plant feeders

## Parasitoids of *Halyomorpha halys* (Heteroptera: Pentatomidae) in Bulgaria

Mariya Hristozova

Agricultural University – Plovdiv, Bulgaria

\*Corresponding author: [mariya\\_valerieva@abv.bg](mailto:mariya_valerieva@abv.bg)

**Abstract:** The brown marmorated stink bug (*Halyomorpha halys*) is an invasive species causing serious damage on many agricultural crops in Bulgaria. The aim of this study was to establish which of the native parasitoid species would attack the eggs of *Halyomorpha halys* and to evaluate the rate of parasitism at field conditions. Egg masses were collected from a number of agricultural crops in the region of Plovdiv and reared under laboratory conditions until hatching. A total of five egg parasitoids emerged from the collected eggs and were identified as *Trissolcus cultratus*, *Trissolcus basalis*, *Anastatus bifasciatus*, *Ooencyrtus telenomicida*, and *Ooencyrtus sp.* The rate of parasitism in 2021 was 10.57% in June, 21.17% in July and 17.57% August.

**Keywords:** *Halyomorpha halys*, egg parasitoids, Scelionidae, Eupelmidae, Encyrtidae

### Subgroup 3 - Plant feeders

## ***Tuta absoluta* egg density and distribution on tomato plants affect the efficacy of mirid predators**

**Sofia A. Dervisoglou<sup>\*1</sup>, Dionysios Ch. Perdikis<sup>1</sup>, Nikos E. Papanikolaou<sup>1,2,†</sup>, Argyro A. Fantinou<sup>3</sup>**

<sup>1</sup>Laboratory of Agricultural Zoology and Entomology, Department of Crop Science, Agricultural University of Athens, Iera Odos 75, 11855, Athens, Greece

<sup>2</sup>Department of Plant Protection Products and Biocides, Hellenic Ministry of Rural Development and Food, Athens, Greece

<sup>3</sup>Laboratory of Ecology and Environmental Science, Department of Crop Science, Agricultural University of Athens, Iera Odos 75, 11855, Athens, Greece

\* Corresponding author: Sofia A. Dervisoglou, e-mail: [sofi.derv@gmail.com](mailto:sofi.derv@gmail.com)

**Abstract:** *Tuta absoluta* is the most serious pest of tomato crops. Over reliance to chemical control has led to its resistance development, adverse effects on human health and deterioration of the product quality. Alternatively, the predators *Macrolophus pygmaeus* (Mp) and *Nesidiocoris tenuis* (Nt) (Hemiptera: Miridae) are effective in its control. However, their efficacy under variable egg densities and distribution patterns on tomato plants has been little searched. In this study the predation rate of single predators (Mp/Nt), conspecific (2Mp/2Nt) or heterospecific pairs (MpNt) was recorded on tomato plants where eggs of *T. absoluta* had been placed on the respective positions that *T. absoluta* females laid their eggs when used at different densities (1, 2, 3, 10 females/plant). Our results showed that the number of eggs laid per *T. absoluta* female was highest when used singly and the eggs were mainly detected on 3<sup>rd</sup> and 4<sup>th</sup> leaves counting from the top. Single *M. pygmaeus* showed a higher predatory efficacy than *N. tenuis* when 14 and 30 eggs were used per plant. Conspecifics of *M. pygmaeus* were more effective than *N. tenuis* at 17 and 110 eggs. A pair was as effective as a single predator at the 14, 17 and in many cases at the 30 and 110 egg densities. *M. pygmaeus* (singly or in combination) may have a higher efficacy than *N. tenuis* at lower and higher *T. absoluta* pest pressure levels. This information is valuable for more effective use of these predators in the management of *T. absoluta*.

**Keywords** *Tuta absoluta*, egg distribution, predatory mirids, biological control, predation rate



## Subgroup 3 - Plant feeders

### Aphids flight activity on wheat

Ivana Lalićević\*, Anđa Radonjić

Department of Phytomedicine, Faculty of Agriculture, University of Belgrade, Nemanjina 6, 11081,  
Zemun, Serbia

\* Corresponding author: [Ivanica1992@gmail.com](mailto:Ivanica1992@gmail.com)

**Abstract:** Aphids (Aphididae, Hemiptera) cause significant economic loss on wheat by direct feeding and by vectoring the viruses. The aim of this study was the monitoring of aphid flight activities and species composition in wheat during the whole season, from beginning of November to middle of June. Study was conducted by yellow water traps in PSS institute Tamiš in Pančevo during autumn 2020/spring 2021. Traps were placed into 18 wheat plots of 60m<sup>2</sup>. Samples were taken once per week. Aphids were identified using a stereoscopic microscope and keys for identification of winged aphids. 42 different taxa were collected; 35 during the autumn and 23 in spring and total of 967 specimens of aphids. The maximum of aphid flight activities was in the middle of November and the end of May. During November the most abundant species was *Myzus persicae* and wheat aphids *Rhopalosiphum padi* and *Sitobion avenae*. During spring the most abundant were specimens of the genus *Aphis*. *Myzus persicae* and *Acyrtosiphon pisum* were presented in high number. Species were also found: *Rhopalosiphum maidis*, *Brachycaudus helichrysi*, *Brachycaudus cardui*, *Dysaphis plantaginea*, *Panaphis juglandis*, *Hyperomyzus lactucae*, *Lipaphis erysimi*, *Wahlgreniella ossiannilssoni*, *Metopolophium dirhodum*, *Phyllaphis fagi*, *Aulacorthum solani*, *Rhopalomyzus poae*, *Schizaphis graminum*, *Phorodon humuli*, *Cinara costata*, *Cavariella aegopodii*, *Myzus cerasi*, *Myzus ascalanicus*, *Macrosiphum rosae*, *Macrosiphum euphorbiae*, *Therioaphis trifolii*. Some of the collected specimens were identified to the genera level: *Anoecia* sp., *Hyadaphis* sp., *Brachycaudus* sp., *Cavariella* sp., *Erisoma* sp., *Hyperomyzus* sp., *Pemphigus* sp., *Myzocallis* sp., *Tetraneura* sp., *Uroleucon* sp. *Dysaphis* sp., etc.

**Keywords:** Aphids, wheat, flight activity, autumn, spring

### Subgroup 3 - Plant feeders

## Natural enemies of *Parthenolecanium corni* (Bouché) (Hemiptera: Coccidae) on grapevine in Neštin, Serbia

Marija Milošević<sup>1\*</sup>, Aleksandar Stojanović<sup>2</sup>, Draga Graora<sup>1</sup>

<sup>1</sup>University of Belgrade - Faculty of agriculture, Belgrade, Serbia

<sup>2</sup>Natural history museum, Belgrade, Serbia

\* Corresponding author: [marija.simonovic@agrif.bg.ac.rs](mailto:marija.simonovic@agrif.bg.ac.rs)

**Abstract:** Natural enemies are valuable part of every entomofauna, especially in agroecosystems. Many of them are used as biological control agents for controlling different pests, including scale insects (Coccidea). With alternative ways for controlling pests in agriculture being a rising trend in plant protection, aim of this study was to determine natural enemies of *Parthenolecanium corni* (Bouché) and their abundance in vineyard with and without insecticide application. The research was conducted during 2019 in commercial vineyard in Neštin locality (northwest Serbia). Two areas of 0.5 ha were marked for this research. In one, there was no insecticide application while in the other insecticides were applied to control other grapevine pests. Examination of vineyard for the presence of entomophagous insects was done twice a month starting from March to October. Twenty randomly chosen vines were examined in both areas. Fourteen natural enemies species were registered, seven predators and seven parasitoids. In untreated area all fourteen species were recorded. From predators, there were registered three species from Coccinellidae (Coleoptera) family and one from each of the following families: Anthribidae (Coleoptera), Forficulidae (Dermaptera), Chrysopidae and Hemerobriidae (Neuroptera). From parasitoids, six species from Encyrtidae and one species from Aphelinidae families were registered. The most abundant natural enemy was *B. longipennis* (48.61%). In treated area total of six entomophagous insects were collected. From predators, one species was registered from each of the families Coccinellidae and Forficulidae. From parasitoids there were recorded three species from Encyrtidae and one from Aphelinidae families. The most abundant natural enemy in treated area was *B. longipennis* (68.09%).

**Keywords:** Coccidae, grapevine, parasitoids, predators, abundance

### Subgroup 3 - Plant feeders

## Some important invasive species in agricultural and urban environments of Serbia

Mihaela Kavran\*, Aleksandra Konjević, Aleksandra Ignjatović Čupina, Miloš Petrović, Dušan Petrić

University of Novi Sad, Faculty of Agriculture, Trg Dositeja Obradovića 8, 21 000 Novi Sad

\* Corresponding author: [mihaela.kavran@polj.edu.rs](mailto:mihaela.kavran@polj.edu.rs)

**Abstract:** Invasive insect species are those that have been introduced to a new natural environment from other parts of the world. These species usually spread as stowaways during transportation, or as a contaminant with commercial goods. Not all alien insect species successfully establish themselves in a new environment. Species which adapted to the moderate climate of Serbia successfully established their populations in the environment with favourable conditions in combination with the absence of natural enemies. This allowed them to spread and reproduce excessively, competing with native species for habitat and resources. Some invasive species may transmit parasites and disease causing agents endangering the health of cultivated plants, native wildlife species, domestic animals and humans.

Invasion of some important insect species in Serbia will be described: *Nezara viridula*, *Halyomorpha halys*, *Bemisia tabaci*, *Aedes albopictus* and *Aedes japonicus*. The first three species are significant agricultural pests, while mosquitoes, but also stink bugs are molestandants of high importance in urban areas. *Nezara viridula* was detected in Serbia in 2008, while *H. halys* invaded Serbia in 2015. These stink bugs are polyphagous, and endanger many agricultural plants. *Bemisia tabaci* is not yet established in Serbia, but its presence was documented several times in greenhouses (2016, 2022). The two invasive mosquito species *Ae. albopictus* and *Ae. japonicus*, both confirmed vectors of several pathogenic agents causing human diseases, were recorded in Serbia for the first time in 2009 and 2018, respectively. Considering the medical importance, *Ae. albopictus* is declared as the most dangerous invasive species in Europe.

**Keywords:** invasive insects, stinkbugs, mosquitoes, urban areas

## Subgroup 3 - Plant feeders

### Sustainable control of insect pests in Mediterranean agroecosystems

**Gianvito Ragone**

*Università degli Studi di Bari - Aldo Moro, via Amendola 165/A, 70126 – Bari, Italy*

\* Corresponding author: [gianvitoragone1@gmail.com](mailto:gianvitoragone1@gmail.com)

**Abstract:** Agricultural expansion and intensification with the consequent loss of seminatural habitats and increased application of agrochemicals, are considered major drivers of biodiversity loss. The reduction in the abundance and diversity of natural enemies inhabiting agricultural landscapes can in fact lead to an ineffective natural control of pests, with potential negative effects on production. This study focuses on three important habitats in Mediterranean agroecosystems: vineyards, olive groves and seminatural habitats. The research aims to assess the effect of local habitat and the characteristics of the surrounding landscape on the abundance and diversity of natural enemies and on the control of phytophagous insects. The evaluation will be carried out in twelve landscapes selected along a gradient in seminatural habitats. Within each landscape we will monitor from May to November the three habitats using different methodologies: phytophagous insects will be sampled with sticky traps and observation of the damage on grapes and olives, whereas natural enemies will be collected through pitfall traps, sweep netting and beating net sampling. Seed cards and dummy caterpillar will be exposed in order to measure predation rates in the field. We expect i) seminatural habitats to be an important source of natural enemies, ii) extensive local management in vineyards and olive groves and high cover of natural habitats in the surrounding landscape to generally support the communities of natural enemies, favouring their potential for biological control. However, iii) some pest species may also benefit from extensive management and the presence of seminatural habitats, potentially generating trade-offs.

**Keywords:** vineyard, olive, landscape, dummy caterpillar, seed cards

### Subgroup 3 - Plant feeders

## Identification of *Trichogramma* egg parasitoids of the European corn borer *Ostrinia nubilalis* (Hübner, 1796) in Vojvodina (Serbia)

Aleksandar Ivezić\*, Aleksandra Ignjatović Čupina

Biosense Institute, University of Novi Sad, Dr Zorana Đinđića 1, 21000 Novi Sad, Serbia

\* Corresponding author: [aleksandarivezic@yahoo.com](mailto:aleksandarivezic@yahoo.com)

**Abstract:** Maize (*Zea mays* Linnaeus, 1753) is the most important cereal crop in Serbia, while the European corn borer (ECB) *Ostrinia nubilalis* (Hübner, 1796) (Lepidoptera: Crambidae) is the major pest of this crop. Regular monitoring of ECB in the province of Vojvodina demonstrated a high percentage of parasitized egg clusters of this pest. Preliminary identification of the egg parasitoids based on morphological traits of emerged adults revealed that they belong to the genus *Trichogramma* Westwood, 1833 (Hymenoptera, Trichogrammatidae). Considering that the fauna of *Trichogramma* wasps has not yet been enough explored in Serbia, the objective of this study was to identify the autochthonous species of *Trichogramma* parasitoids of ECB present in the region of Vojvodina by using sequences of ribosomal DNA. The research conducted in Vojvodina during 2016 and 2017 revealed presence of two *Trichogramma* species: *Trichogramma brassicae* and *Trichogramma evanescens*. Intraspecific variation of samples of wasps was assessed by amplifying the mitochondrial cytochrome oxidase I (COI) gene. The results showed the presence of 4 haplotypes of *T. brassicae* in local populations, and a single haplotype of *Trichogramma evanescens*. Wasps were also tested for the presence of the endosymbiotic bacteria *Wolbachia* using *Wolbachia*-specific 16S and *wsp* primers and the results demonstrated the presence of the bacteria in the population of *T. brassicae* in its dominant haplotype (TbSERB-1). Understanding of the diversity and biology of autochthonous *Trichogramma* species is a key step in the potential commercial implementation of these egg parasitoid wasps in the protection of corn fields in Serbia.

**Keywords:** beneficial insects, biological control, *Ostrinia nubilalis*, *Trichogramma* spp, maize

### Subgroup 3 - Plant feeders

## Fauna of insects from the suborder Sternorrhyncha in citrus orchards in Dubrovačko -neretvanska County

Vjekoslav MARKOTIĆ<sup>1\*</sup>, Renata BAŽOK<sup>2</sup>, Tatjana MASTEN MILEK<sup>3</sup>, Mladen ŠIMALA<sup>1</sup>

<sup>1</sup>Croatian Agency for Agriculture and Food, Centre for Plant Protection, Gorice 68b, 10000 Zagreb,  
Croatia

<sup>2</sup>University of Zagreb Faculty of Agriculture, Department of Agricultural Zoology, Svetošimunska cesta  
25, 10000 Zagreb, Croatia

<sup>3</sup>Public Institution Green Ring, 151. samoborske brigade HV 1, 10430 Samobor

\*Corresponding author: [vjekoslav.markotic@hapih.hr](mailto:vjekoslav.markotic@hapih.hr)

**Abstract:** Earlier faunistic studies of insects of the suborder Sternorrhyncha (Hemiptera) in Croatia included surveys in citrus-growing areas of plant species of the genera *Citrus* and *Poncirus*, but only on a smaller scale. Considering the expansion of trade in planting material in the last decade and the constant findings of quarantine pests both in Mediterranean countries and in Croatia, the aim of this study is to conduct an inventory of insects of the suborder Sternorrhyncha in Dubrovačko-neretvanska County, the most important citrus growing area in Croatia. For six years (2015-2020), faunistic surveys were conducted in citrus orchards, backyards and nurseries in Dubrovačko-neretvanska County. During this period, ten plant species of the genera *Citrus* and *Poncirus* were studied at a total of 171 sites from April to November. Cenological characteristics of the identified insects were established. A total of 17 species belonging to the suborder Sternorrhyncha were found. Significantly more species were found in commercial orchards compared to backyards and nurseries. The whitefly *Dialeurodes citri*, the aphids *Aphis gossypii* and *A. spiraecola*, and the scale insects *Coccus pseudomagnoliarum* and *Aonidiella aurantii* were classified as eudominant. The quarantine species *Aleurocanthus spiniferus* was found in two distant localities, so its eradication is no longer possible due to the current extent of infection. The regulated non-quarantine species *Parabemisia myricae* was also found in two localities and no primary damage to citrus trees or symptoms of infection with the Citrus Chlorotic Dwarf virus transmitted by this species were detected during the surveys.

**Keywords:** cenological characteristics, citrus, insect pests, entomofauna



## Subgroup 4 - Plant Pathology

### Multi-omics approaches to explore phenotypic and genetic diversity in the brown rot fungal pathogens *Monilinia fructicola*, *Monilinia laxa* and *Monilinia fructigena*

Rita Milvia De Miccolis Angelini<sup>1</sup>, Celeste Raguseo<sup>1</sup>, Lucia Landi<sup>2</sup>, Caterina Rotolo<sup>1</sup>, Palma Rosa Rotondo<sup>1</sup>, Donato Gerin<sup>1</sup>, Gianfranco Romanazzi<sup>2</sup>, Stefania Pollastro<sup>1</sup>, Francesco Faretra<sup>1\*</sup>

<sup>1</sup>Department of Soil, Plant and Food Sciences, University of Bari Aldo Moro, Bari, Italy

<sup>2</sup>Department of Agricultural, Food and Environmental Sciences, Marche Polytechnic University, Ancona, Italy

Corresponding author: [francesco.faretra@uniba.it](mailto:francesco.faretra@uniba.it)

**Abstract:** *Monilinia fructicola*, *Monilinia fructigena* and *Monilinia laxa* are the main causal agents of brown rot and blossom blight, among the most important diseases on pome and stone fruits, worldwide responsible for heavy losses both in the field and in postharvest. Following the introduction of *M. fructicola* into Europe in 2001, the pathogen spread widely and rapidly, and became prevalent over the endogenous species *M. laxa* and *M. fructigena*. To improve knowledge on the biology, evolutionary history, and diversity of these important pathogens, high-quality reference genomes and transcriptomes of *M. fructicola* strain Mfrc123, *M. fructigena* strain Mfrg269 and *M. laxa* strain Mlax316 were *de novo* assembled and compared. In particular we explored: (i) phylogenomic and synthenic relationships between the *Monilinia* genomes and those of the closely related species within *Sclerotiniaceae*, *Botrytis cinerea* and *Sclerotinia sclerotiorum*; (ii) the abundance and evolutionary dynamics of transposable element; (iii) the genetic basis of mating type and the genome-wide occurrence and extent of Repeat-Induced Point (RIP) mutations; and (iv) common and species-specific effectors, carbohydrate-active enzymes and secondary metabolite gene clusters with some differences that might explain host plants and organ preferences distinguishing the three *Monilinia* species. Moreover, a metagenomic approach was used to investigate the mycovirome of *M. fructicola* in a worldwide collection of isolates from different hosts and revealed a great abundance and variety of mycoviruses infecting the fungus. A total of 32 positive-sense single-stranded (ss)RNA viruses and a new putative ssDNA mycovirus were identified and characterized.

**Keywords:** Next-generation sequencing, *de novo* assembly, comparative genomics, transcriptomics, metagenomics, mycovirome

## Subgroup 4 - Plant Pathology

### Biocontrol activity of epiphytic yeasts against *Penicillium expansum* on apple fruits

Tatjana Dudaš<sup>1\*</sup>, Dragana Budakov<sup>1</sup>, Simona Sanzani<sup>2</sup>, Mila Grahovac<sup>1</sup>, Marta Loc<sup>1</sup>, Vera Stojšin<sup>1</sup>, Antonio Ippolito<sup>2</sup>

<sup>1</sup>University of Novi Sad, Faculty of Agriculture, Department of Plant and Environmental Protection

<sup>2</sup>Dipartimento di Scienze del Suolo, della Pianta e degli Alimenti, Università degli Studi di Bari Aldo Moro, Via Amendola 165/A, 70126 Bari, Italy

\* Corresponding author: [tatjana.dudas@polj.uns.ac.rs](mailto:tatjana.dudas@polj.uns.ac.rs)

**Abstract:** *Penicillium expansum* causes apple blue mold resulting in economic losses and threatening human health due to patulin production. Synthetic fungicides are used for the control of the pathogen, but new strategies are necessary for sustainable production, and epiphytic yeasts as biocontrol agents are a possible alternative. Yeast strains were collected from the surface of healthy untreated apple fruits (Dudaš et al., 2022). Biocontrol activity of 17 yeast strains was tested on apple fruits following the method by Sanzani et al. (2021). Four wounds (3x3 mm) were made on Golden Delicious apple fruits and treated with 10 µl of yeast suspension ( $10^7$  cells/ml) and with 10 µl of pathogen suspension ( $10^4$  cells/ml). In fungicide treatment, wounds were treated with 10 µl of thiophanate methyl (0.05 %) instead of yeast. Untreated fruits served as the negative control, while fruits treated with sterile distilled water and afterwards with the pathogen were positive control. Four fruits were used per treatment. Lesion diameters were measured after 9 days at 20 °C. All yeast strains except one caused statistically significant reduction of lesion diameters. Treatment with fungicide resulted in an average lesion diameter of 38.7 mm and showed no statistically significant difference with the positive control (39.1 mm) as well as one yeast strain (38.9 mm). One yeast strain showed the strongest biocontrol activity, with an average lesion diameter of 4.5 mm. Further insights into the biocontrol mode of action of the most effective yeast strain will be conducted, as well as larger-scale trials to assess possible practical field applications.

**Keywords:** *Penicillium expansum*, apple blue mold, biocontrol, epiphytic yeasts

Dudaš, T., Budakov, D., Loc, M., Grahovac, M., Stojšin, V. (2022): Biocontrol activity of antagonistic yeasts against *Penicillium expansum* – major patulin producer in apples. The World Mycotoxin Forum - 13 th conference WMF Meets Italy. 16-18 May, 2022, Parma, Italy. p.138.

## Subgroup 4 - Plant Pathology

Sanzani, S. M., Sgaramella, M., Mosca, S., Solfrizzo, M., Ippolito, A. (2021): Control of *Penicillium expansum* by an Epiphytic Basidiomycetous Yeast. *Horticulturae* **2021**, 7, 473.  
<https://doi.org/10.3390/horticulturae7110473>

## Subgroup 4 - Plant Pathology

### Overview of the use of GWAS to identify loci for resistance to apple scab caused by *Venturia Inaequalis*

Konjić Almira\*, Arnela Okić, Jasmin Grahić, Fuad Gaši

<sup>1</sup>Univerzitet u Sarajevu, Poljoprivredno-prehrambeni fakultet, Sarajevo, Bosna i Hercegovina

\* Corresponding author: [a.konjic@ppf.unsa.ba](mailto:a.konjic@ppf.unsa.ba)

**Abstract:** Apple scab caused by fungus *Venturia inaequalis* is the most devastating disease of the cultivated apple (*Malus domestica*). It is the main limiting factor of a successful production due to losses in fruit quality and yield, causing farmers to treat orchards with fungicides multiple times per season. In this study we will explore the use of Genome Wide Association Study (GWAS) on single nucleotide polymorphism (SNP) to identify genomic regions associated with resistance to apple scab. Obtained results would help us to better understand the resistance to *Venturia inaequalis* in apple and show significant information for future resistance breeding programs.

**Keywords:** SNP, GWAS, Apple scab, disease resistance, resistance breeding

## Subgroup 4 - Plant Pathology

### Endophytic bacteria from olive drupes as biological control and induced resistance agents of *Colletotrichum acutatum*

Anastasia Papageorgiou, Maria Varveri, Dimitrios Tsitsigiannis\*

Laboratory of Plant Pathology, Department of Crop Science, Agricultural University of Athens, Greece

\* Corresponding author: [dimtsi@aua.gr](mailto:dimtsi@aua.gr)

**Abstract:** Olive anthracnose (OA) is one of the most important diseases affecting olives worldwide. Causal agents of OA are fungi belonging to the genus *Colletotrichum*, while the predominant one is *C. acutatum* species complex. Due to the fungicide withdrawals and the risk of pathogen resistance, finding more sustainable control measures of the disease becomes extremely necessary. The aim of this study is to investigate whether endophytic microorganisms from olive drupes are able to reduce OA disease severity and induce plant's defence mechanisms. Hence, 210 endophytic bacteria were isolated and tested for their antagonistic effect on *C. acutatum* in multiple and dual culture essays. Bacterial isolations that demonstrated satisfactory inhibition of the mycelial growth, were further tested for their effect on the disease severity and conidial production of the fungus, in olive drupes of cv. Kalamon. The most effective isolates were evaluated in olive trees, artificially inoculated by the pathogen, where the bacterium П8 (*Serratia* spp.) caused a reduction of the disease severity by approximately 90%. Those isolates were, also, applied on young olive seedlings in greenhouse conditions in order to assess their effect on olive plant defence mechanisms. Total RNA was extracted from leaves, and the expression of ten different defence genes was evaluated by RT-qPCR. All four tested bacterial strains caused increased expression of genes associated with Pathogenesis-related proteins, while genes involved in biosynthetic pathways of phenylpropanoids, salicylic and jasmonic acid were induced by the presence of strains K13 (*Bacillus methylotrophicus*), B1 (*B. amyloliquefaciens*) and П8 (*Serratia* sp.).

**Keywords:** *C. acutatum*, defence genes, bacteria, plant- resistance inducers, disease severity

## Subgroup 4 - Plant Pathology

### The volatile organic compounds of the biocontrol agent *Bacillus velezensis* K165 activate plant defenses against *Verticillium dahliae*

Eirini Poulaki<sup>1\*</sup>, Ioannis Stringlis<sup>2</sup>, Marinos Xagoraris<sup>3</sup>, Christos Pappas<sup>3</sup>, Sotiris Tjamos<sup>1\*</sup>

<sup>1</sup> Laboratory of Phytopathology, <sup>3</sup> Laboratory of General Chemistry Agricultural University of Athens, Iera Odos 75, 11855, Athens, Greece, <sup>2</sup> Plant-Microbe Interactions, Department of Biology, Utrecht University, The Netherlands

\* Corresponding author: [poulakie@gmail.com](mailto:poulakie@gmail.com)

**Abstract:** *Verticillium dahliae* is one of the most devastating soilborne pathogens worldwide. Due to the unavailability of effective chemical strategies and the lack of resistant plant cultivars, it is urgent to develop innovative disease management strategies. In this respect, the use of the biocontrol strain *Bacillus velezensis* K165 represents a promising approach. Here, we examined the biocontrol efficacy of the volatile organic compounds (VOCs) produced by K165 against *V. dahliae* in *Arabidopsis thaliana*. Initially, it was found that K165 VOCs-exposed *A. thaliana* plants displayed reduced Verticillium wilt symptoms compared to control plants. Subsequently, analysis of RNAseq data and pathogenicity experiments employing *A. thaliana* mutants showed the positive interplay between K165 VOCs and the chitin perception mechanism of plants. The GC/MS analysis of the K165 VOCs profile revealed acetoin as the major chemical compound emitted by K165. Acetoin did not reduce the growth of *V. dahliae* *in vitro*. Nevertheless, acetoin-exposed *A. thaliana* plants displayed reduced Verticillium disease severity compared to controls. Therefore, the K165 VOCs seem to activate the defense system of plants and contribute to the biocontrol activity of K165 against *V. dahliae*.

**Keywords:** acetoin, biological control, chitin, induced systemic resistance, soilborne pathogens





## Subgroup 4 - Plant Pathology

### Botryosphaeriaceae as postharvest pathogens of apple fruit

Nina Vučković\*, Ivana Vico, Nataša Duduk

University of Belgrade - Faculty of Agriculture, Nemanjina 6, Belgrade, Serbia

\* Corresponding author: [ninaradulovic@hotmail.com](mailto:ninaradulovic@hotmail.com)

**Abstract:** Species of the family Botryosphaeriaceae are important pathogens of woody plants, including apple fruit trees, known to cause stem and branch cankers, dieback, and pre- and postharvest fruit rot. In the survey conducted from 2016 to 2018, several Botryosphaeriaceae species were isolated from apple fruit with symptoms of brown rot collected from different storages (controlled atmosphere, normal atmosphere, and cellar) and orchards during harvest. The obtained isolates, whose pathogenicity was confirmed on apple fruit, were identified and characterized based on multilocus phylogeny and anamorph morphology. Five species from the family Botryosphaeriaceae were revealed as postharvest pathogens of apple fruit in Serbia. *Diplodia seriata*, followed by *Botryosphaeria dothidea*, were the most prevalent species, both of which were previously recorded on apple fruit in Serbia. *Diplodia bulgarica*, *Diplodia sapinea* and *Neofusicoccum yunnanense* represent first reports on this host in the country. It has been noticed in this study that rot symptoms caused by these species can occur during harvest but are more frequent after harvest following unfavourable storage conditions. This may suggest that Botryosphaeriaceae species persist endophytically, as latent pathogens, in apple fruit and that warmer storage conditions favour disease development. The results of this study imply that species of the family Botryosphaeriaceae represent an increasing threat to successful apple production due to their endophytic nature, pathogenic potential that comes to full expression under plant stress conditions such as global climate change, as well as the ability to cause other diseases on their hosts.

**Keywords:** *Botryosphaeria*, *Diplodia*, *Neofusicoccum*, multilocus phylogeny, apple fruit decay

## Subgroup 4 - Plant Pathology

### Antagonistic activity of endophytic *Bacillus* strains against phytopathogenic fungi. Mode of action studies and the strain compatibility issue

Polina Tsalgaidou<sup>1,2</sup>, Eirini Evangelia Thomloudi<sup>1</sup>, Eirini Baira<sup>3</sup>, Panagiotis Katinakis<sup>1</sup>, Anastasia Venieraki<sup>4\*</sup>

<sup>1</sup>Laboratory of General and Agricultural Microbiology, Crop Science Department, Agricultural University of Athens, Iera Odos 75, 11855 Athens, Greece.

<sup>2</sup> Department of Agriculture, University of the Peloponnese, Antikalamos, Messinia, Greece.

<sup>3</sup>Laboratory of Toxicological Control of Pesticides, Scientific Directorate of Pesticides' Control and Phyto-pharmacy, Benaki Phytopathological Institute (BPI), Kifissia, 14561 Athens, Greece.

<sup>4</sup>Laboratory of Plant Pathology, Crop Science Department, Agricultural University of Athens, Iera Odos 75, 11855 Athens, Greece.

\* Corresponding author: [venieraki@aua.gr](mailto:venieraki@aua.gr)

**Abstract:** The effects of agrochemical overuse on environmental and human health highlighted the necessity of improving and developing biological approaches to achieve increase of crop yield and decrease of losses. In this study, we isolated and analysed for their beneficial characteristics endophytic bacterial strains of the genus *Bacillus* with antagonistic effect against several phytopathogens. We tested their benefits under *in vitro* and *ex vivo* conditions. All *Bacillus* strains studied successfully colonized the plant tissues and competed with the phytopathogenic fungi for nutrients and space. We applied an integrative approach coupling genome mining and metabolic profiling to decipher the potential of these endophytes as biological control agents against pathogenic fungi. The chemical analysis of their extract indicated that these *Bacillus* endophytes were able to secrete several bioactive secondary metabolites. The extensive metabolomic (HPLC-HRMS) and whole genome analysis demonstrated the ability of these endophytes to biosynthesize and secrete a significant number of bioactive secondary metabolites (fengycin, surfactin, iturin, mojavensin, bacilysin etc.) and lytic enzymes (chitinase, cellulase, protease, etc.). All of these compounds determine their competitiveness, colonization and survival ability, the compatibility with each other and the induction of induced systemic resistance (ISR). Also, we investigate their ability for fungal growth suppression either as individual strains or as bacterial mixtures. The selected strains were paired for the development of mixtures after assays regarding their *in vitro* compatibility, combining strains with different abilities. Formulation of compatible bacterial strains resulted in some successful mixtures with strong biological control activity.

**Keywords:** Biocontrol, endophytic bacteria, biostimulants, secreted metabolites, strain compatibility

## Subgroup 4 - Plant Pathology

### Antifungal activity of peppermint, sweet basil, and sage essential oils on plant pathogenic fungi

Saud Hamidović\*, Fejzo Bašić, Sanel Haseljić, Teofil Gavrić, Arnela Okić, Berina Imamović, Nedžad Karić

<sup>1</sup>Univerzitet u Sarajevu, Poljoprivredno-prehrambeni fakultet, Sarajevo, Bosna i Hercegovina

\* Corresponding author: [s.hamidovic@ppf.unsa.ba](mailto:s.hamidovic@ppf.unsa.ba)

**Abstract:** The application of pesticides is still a common practice to protect plants, fresh fruits, and vegetables, which aims to ensure adequate yield and product quality. However, some of the chemical compounds used can have an adverse effect on human health, enter the food chain, and some microorganisms become resistant to the chemicals used. Due to these concerns, there is an increasing need to find new, more effective, and safe alternative protective agents of natural origin. The aim of this work was to investigate the antifungal effect of peppermint (*Mentha piperita*), basil (*Ocimum basilicum*), and sage (*Salvia officinalis*) oils on the growth of fungi from the genus *Fusarium* sp. and *Aspergillus* sp. The oil was extracted from dried plant stems and leaves by hydrodistillation, using an adapted Neo Clevenger apparatus, according to ISO-standard 857-1984 (E). The antifungal activity was determined by the test-diffusion method on a PDA substrate. Paper discs with a diameter of 6 mm were impregnated with 15 ml of selected oils and placed on PDA with inoculated pure cultures of the fungi. The incubation was performed at 22°C. The inhibition zones were measured after the third and sixth day of incubation. In the control variant, distilled water was used. The results showed a reduction in the growth of the tested fungi after the incubation period, indicating a potential use of the analyzed oils in protection against fungi from the genus *Fusarium* sp. and *Aspergillus* sp.

**Keywords:** antifungal activity, essential oils, *Fusarium* sp., *Aspergillus* sp.

## Subgroup 4 - Plant Pathology

### Physiological response in winter wheat varieties depending on *Fusarium* infection and N fertilization

Magdalena Matic\*

Faculty of Agrobiotechnical Sciences Osijek, Josip Juraj Strossmayer University of Osijek,  
Vladimira Preloga 1, Osijek, Croatia

\* Corresponding author: [matimc@fazos.hr](mailto:matimc@fazos.hr)

**Abstract:** Wheat (*Triticum aestivum* L.) is one of the most important cereal crops in the world. Under field conditions, wheat is exposed to both abiotic and biotic stresses that have a great impact on its yield and quality. Inadequate nitrogen fertilization represents an abiotic stress factor that causes numerous physiological and metabolic changes in wheat tissues. An important biotic stress factor in wheat production is the occurrence of *Fusarium* head blight (FHB), a disease caused by phytopathogenic species of the genus *Fusarium*. These stressors enhance the rapid accumulation of reactive oxygen species (ROS) and the occurrence of oxidative stress in plant tissues. However, plants have developed enzymatic and non-enzymatic antioxidant defense mechanisms to control the excessive ROS levels. For the purpose of this study, the field experiment was set up during two growing seasons in the experimental field of the Agricultural Institute Osijek. It was set-up in a split-split-plot design as a complete randomized block in two replications, as a factorial experiment with three main factors: variety, N fertilization levels, and *Fusarium* infection. Biomarkers of oxidative stress and antioxidant response were determined in wheat samples. The aim of this study was to determine the variety-specific oxidative and antioxidant response of wheat to different nitrogen fertilization and FHB and to determine potential biochemical mechanisms of wheat resistance.

**Keywords:** wheat, nitrogen, *Fusarium* spp., oxidative stress, antioxidant response

## Subgroup 4 - Plant Pathology

### Grafting as a sustainable approach in response to infections transmitted by arthropods

**M. Mastrochirico, Roberta Spanò, Tiziana Mascia\***

*Università degli Studi di Bari, Dipartimento di Scienze del Suolo, della pianta e degli Alimenti,  
via Amendola 165/A, 70125, Italy.*

\* Corresponding author: [tiziana.mascia@uniba.it](mailto:tiziana.mascia@uniba.it)

**Abstract:** Grafting is an agronomic technique used as an integrated strategy for the management of abiotic and biotic stresses like bacteria, viruses and fungi in a sustainable and environmentally friendly approach. Grafted tomato plants could also tolerate infections of air-borne viruses like the CiPz Sw5-breaking strain of *tomato spotted wilt virus* (TSWV-CiPz), strains of *cucumber mosaic virus* supporting stunting or necrotic satellite RNAs (CMV-TTS and CMV-77, respectively) and of a recombinant strain of *potato virus Y* (PVY<sup>C-to</sup>), necrotic to tomato. The graft promotes the transfer of tolerance from the rootstock to the scion in susceptible tomato F1 UC82 (UC) varieties on the tolerant tomato local ecotype, denoted Manduria (Ma). The grafting technique reduced the differentially expressed genes (DEGs) in the UC and Ma plants by approximately 54% upon the infection with PVY<sup>C-to</sup>. Moreover, viral infection modulated the up-regulation of specific differentially expressed genes (DEGs) that were significantly expressed in susceptible variety UC and in UC and Ma self- or hetero-grafted plants. It is proposed to use the same type of approach to obtain useful levels of tolerance to *tomato leaf curl New Delhi virus* (ToLCNDV) for cucurbits. ToLCNDV is an emerging begomovirus listed in the EPPO Alert-list 2, present in the mediterranean area, including South-Italy. Twenty-one local cucurbit ecotypes were screened to assess tolerance levels against mechanical transmission of ToLCNDV. The results will lead to the identification of tolerant cucurbit ecotypes to use as potential rootstocks to achieve adequate levels of tolerance against the virus in commercial cucurbit varieties.

**Keywords:** vegetable grafting, plant viruses, disease tolerance, RNAi, sustainability



## Subgroup 4 - Plant Pathology

### Cold Plasma as a new technology in the sustainable management of plant diseases

Palma Rosa Rotondo<sup>1</sup>, Rita Milvia De Miccolis Angelini<sup>1\*</sup>, Paolo Francesco Ambrico<sup>2</sup>, Caterina Rotolo<sup>1</sup>, Donato Gerin<sup>1</sup>, Stefania Pollastro<sup>1</sup>, Francesco Faretra<sup>1</sup>

<sup>1</sup>Department of Soil, Plant and Food Sciences, University of Bari 'Aldo Moro', Bari, Italy

<sup>2</sup>Consiglio Nazionale delle Ricerche, Istituto per la Scienza e la Tecnologia dei Plasmi, Bari, Italy

\*Corresponding author: [ritamilvia.demiccolisangelini@uniba.it](mailto:ritamilvia.demiccolisangelini@uniba.it)

**Abstract:** In the last decade, the need of sustainable, innovative and efficient control methods is a primary goal for the management of plant disease. Non-thermal plasma, also known as cold plasma, is proposed as a green technology for eco-friendly control of phytopathogens on fruits, seeds and plants. It is a partially ionized gas, composed of a mixture of ions, electrons, radicals, reactive oxygen and nitrogen species (RONS) and UV radiation that have a specific role in the decontamination activity carried out directly by air plasma or indirectly by water activated by plasma (PAW). Working conditions at atmospheric pressure and ambient temperature makes this technology suitable for biological application. The *in vitro* efficacy of cold plasma against different fungal and bacterial species was evaluated using different configurations of plasma sources, *i.e.* Surface Dielectric Barrier Discharge (SDBD), Atmospheric Pressure Plasma Jet (APPJ), Volume Dielectric Barrier Discharge (VDBD) and treatment conditions. Albino mutants of *Botrytis cinerea* and *Aspergillus carbonarius* was used to explore the protective role of fungal melanin on sensitivity to plasma exposure, with mutants showing higher sensitivity to treatments compared to the melanized wild-type strains. Efficacy of plasma treatments against pathogenic fungi and bacteria contaminating seeds was evaluated on tomato seeds artificially inoculated with *Cladosporium fulvum* and *Pseudomonas syringae* pv. *tomato* and on barley seeds inoculated with *Fusarium* spp.. Experiments on cherry and citrus fruits demonstrated that the application of plasma may significantly extend their shelf life during postharvest storage by direct microbial inactivation and possible activation of plant defence responses.

**Keywords:** Non-thermal plasma, green technology, phytopathogens, melanin, decontamination



## Subgroup 4 - Plant Pathology

### Response of indigenous tetraploid wheat landraces from Montenegro to infection caused by Barley Yellow Dwarf Virus

Velimirovic Ana<sup>1</sup>, Jovovic Zoran<sup>1</sup>, Perovic Dragan<sup>2</sup>, Mandic Dragan<sup>3</sup>, Mikic Sanja<sup>4</sup>, Przulj Novo<sup>5</sup>

<sup>1</sup>Biotechnical Faculty, University of Montenegro<sup>2</sup>Julius Kühn-Institut Federal Research Institute for Cultivated Plants,<sup>3</sup>Agricultural Institute of Republic of Srpska, Bosnia and Hercegovina, <sup>4</sup>Institute of Field and Vegetable Crops, Novi Sad, Serbia, <sup>5</sup>Faculty of Agriculture, University of Banja Luka

\* Corresponding author: [ana.velimirovic@hotmail.com](mailto:ana.velimirovic@hotmail.com)

**Abstract:** Barley Yellow Dwarf Virus (BYDV) is one of the most economically important viruses in cereals. BYDV causes substantial losses throughout the world wherever wheat, barley, and oats are grown. The yield loss caused by this virus depends on aphid activity, BYDV strain, growth stage at infection, and environmental conditions. Montenegrin collection consists of 85 indigenous tetraploid wheat populations belonging to alternative seasonal types. Twenty spikes of each population were sown in Banjaluka, Bosnia, and Hercegovina during 2020. Sowing was performed in a spike per row system. During the vegetation period, plants originating from five populations were completely destroyed by this virus in the tillering growth stage. This result shows that BYDV did not affect a large fraction of a large part of the Montenegrin tetraploid wheat population (94%). Since the control of the virus itself is limited, the only mechanism of infection control is engagement in the production of resistant or tolerant genotypes. Considering the devastating effect of this virus on the wheat, there is an indication that local populations from Montenegro might be the source of resistance or tolerance mechanisms to BYDV.

**Keywords:** BYDV, tetraploid wheat, landraces, resistance mechanisms

## Subgroup 4 - Plant Pathology

### Studying epidemiology of *Xanthomonas euvesicatoria* strains using molecular genotyping methods

Jelena Menković\*, Aleksa Obradović  
University of Belgrade, Faculty of Agriculture

\* Corresponding author: [jelena.menkovic@agrif.bg.ac.rs](mailto:jelena.menkovic@agrif.bg.ac.rs)

**Abstract:** Bacterial spot, caused by *Xanthomonas euvesicatoria*, is one of the most devastating bacterial diseases of pepper (*Capsicum annuum* L.). The research will include strains originating from different localities and different years of isolation. Newly isolated *X. euvesicatoria* strains, as well as strains stored in the Laboratory Collection, both identified by classical bacteriological and molecular tests, will be subjected to epidemiological studies by using molecular genotyping methods. Genotyping will be performed by analyzing the variability in the number of short repetitive gene sequences at different loci in the bacterial genome (multiple loci variable number of tandem repeat analysis - MLVA), as well as by comparing the genetic similarity of the strains isolated in Serbia with strains from neighboring countries in order to determine level of genetic similarities and distribution of the pathogen population. MLVA is a very popular assay in bacterial genotyping due to its high level of specificity, reproducibility, efficiency and low cost of performance. With this method, it can be determined whether there are similarities between strains from different regions and countries, which facilitates monitoring spread of the pathogen into new areas.

**Keywords:** Pepper, *Xanthomonas euvesicatoria*, MLVA

## Subgroup 4 - Plant Pathology

### Root rot of sugar beet and strawberry caused by *Rhizoctonia* spp. in Serbia

Mira Vojvodić<sup>1\*</sup>, Ivana Vico<sup>1</sup>, Brankica Tanović<sup>2</sup>, Petar Mitrović<sup>3</sup>, Dejan Lazić<sup>4</sup>,  
Karolina Vrandečić<sup>5</sup>, Aleksandra Bulajić<sup>1</sup>

<sup>1</sup> University of Belgrade-Faculty of Agriculture, Nemanjina 6, Belgrade, Serbia

<sup>2</sup> Institute of Pesticides and Environmental Protection, Banatska 31b, Belgrade, Serbia

<sup>3</sup> Institute of Field and Vegetable Crops, Maksima Gorkog 30, Novi Sad, Serbia

<sup>4</sup> Biotech engineering d.o.o., Golsvordijeva 32, Belgrade, Serbia

<sup>5</sup> Faculty of Agrobiotechnological Sciences, Josip Juraj Strossmayer, University of Osijek, Vladimira  
Preloga 1, Osijek, Croatia

\*Corresponding author: [miravojvodic2510@gmail.com](mailto:miravojvodic2510@gmail.com)

**Abstract:** *Rhizoctonia* spp. are economically important soil-borne pathogens infecting a wide range of host plants. Due to the prominent diversity, according to cell nuclear condition, *Rhizoctonia* spp. are classified as multinucleate (teleomorphs: *Thanatephorus* and *Waitea*), binucleate (teleomorphs: *Ceratobasidium* and *Tulasnella*) and uninucleate (teleomorph: *Ceratobasidium*), whereas multinucleate *Rhizoctonia* are further divided into 13 (AG-1 -13) and binucleate *Rhizoctonia* into 23 anastomosis groups (AG-A -W).

At two localities in Serbia, in the vicinity of Belgrade and Vašica (Belgrade City and Srem Districts), sugar beet and strawberry plants expressing prominent root and crown necrosis were sampled and multiple, morphologically uniform isolates were obtained. Two representative isolates originating from sugar beet and strawberry (270-15 and 101-16) were identified based on hyphal branching pattern, nuclear status and conventional AG pairing with known tester isolates. The identification was confirmed by sequencing of ITS region of rDNA and isolate 270-15 from sugar beet was identified as *R. solani* AG-2-2 (Acc. No MK123309) while isolate 101-16 from strawberry proved to be binucleate *Rhizoctonia* sp. AG-A (Acc. No MH517393). Pathogenicity was confirmed after artificial inoculations of strawberry runners and sugar beet seedlings, respectively.

Managing *Rhizoctonia* root rot of sugar beet and strawberry is difficult due to the soil-borne nature and wide host range of the pathogen. There are differences among AGs regarding host range and fungicide sensitivity, so identification of *Rhizoctonia* AG causing root rot of diseased host, including sugar beet and strawberry, is the first step in developing a successful disease management strategy.

**Keywords:** *Rhizoctonia*, root rot, sugar beet, strawberry, identification

## Subgroup 4 - Plant Pathology

### The effect of plant essential oils on *in vitro* growth of *Pectobacterium punjabense*, *Pectobacterium brasiliense* and *Pectobacterium carotovorum*

Marta Loc<sup>1\*</sup>, Dragana Milošević<sup>2</sup>, Milica Aćimović<sup>3</sup>, Tatjana Dudaš<sup>1</sup>, Dragana Budakov<sup>1</sup>, Vera Stojšin<sup>1</sup>, Mila Grahovac<sup>1</sup>

<sup>1</sup> Department of Plant and Environmental Protection, Faculty of Agriculture, University of Novi Sad, Trg Dositeja Obradovića 8, 21000 Novi Sad, Serbia, <sup>2</sup> Laboratory for seed testing, Institute of Field and Vegetable Crops, Novi Sad, Serbia, <sup>3</sup> Department of vegetable and alternative crops, Institute of Field and Vegetable Crops, Novi Sad, Serbia

\*Corresponding author: [marta.loc@polj.edu.rs](mailto:marta.loc@polj.edu.rs)

**Abstract:** *Pectobacterium* spp. have emerged as one of the top 10 phytopathogens affecting plants during production, handling, storage and transportation. These wide distributed pathogens are downgrading the potato industry every year. Many efforts have been made to control bacterial soft rot on potato but with limited success. Essential oils (EOs) and compounds of wild and cultivated plants have been recognized as powerful antibacterial agents. This study evaluated the *in vitro* effect of *Ocimum basilicum*, *Satureja hortensis*, *Origanum vulgare*, *Calluna vulgaris* and *Thymus vulgaris* EOs on *P. punjabense* (MMZCVR2), *P. brasiliense* (MMKC19) and *P. carotovorum* (MMDC11) strains originating from potato from Serbia and two reference strains from Lebanon (DAPP-PG 752) and Brasil (LMG 21370), using vapour diffusion method (Clemente et al., 2016), obtaining concentration of 0.16 µl/ml of air. The results were recorded after 72h of incubation at 26°C. It was found that all tested EOs varied in effect on growth of bacterial strains. Results showed that EO of *S. hortensis* and *C. vulgaris* showed the highest bacterial growth inhibition, significantly higher compared to *Th. vulgaris* EO, while *O. basilicum* and *O. vulgare* EOs showed the lowest growth inhibition, however significantly higher compared to control. The vitality of bacterial colonies was checked by streaking on NA medium and incubation during 48h at 26°C. It was revealed that *S. hortensis* and *C. vulgaris* EOs showed bactericidal effect on all strains, as well as *Th. vulgaris* EO on MMKC19 and both DAPP-PG 752 and LMG 21370. *O. basilicum* and *O. vulgare* EOs showed bacteriostatic effect on all isolates, as well as *Th. vulgaris* EO on MMDC11 and MMZCVR2.

**Keywords:** *Pectobacterium*, potato, blackleg, softrot, essential oils, biocontrol

Clemente, I., Aznar, M., Silva, F. and Nerín, C., 2016. Antimicrobial properties and mode of action of mustard and cinnamon essential oils and their combination against foodborne bacteria. *Innovative Food Science & Emerging Technologies*, 36, pp.26-33.



## Subgroup 4 - Plant Pathology

### Resistance of *Olea europaea* towards *Xylella fastidiosa* subsp. *pauca* ST53: influence of cultural practices and endogenous factors

Mariangela Carlucci<sup>1</sup>, Pierfederico La Notte<sup>2</sup>, Franco Nigro<sup>1\*</sup>

<sup>1</sup>Dipartimento di Scienze del Suolo, della Pianta e degli Alimenti, Università degli Studi di Bari  
- Aldo Moro, via Amendola 165/A, 70126 – Bari, Italy

<sup>2</sup>Istituto per la Protezione Sostenibile delle Piante, Consiglio Nazionale delle Ricerche, via  
Amendola 122/D 70126 - Bari, Italy

\*Corresponding author: [franco.nigro@uniba.it](mailto:franco.nigro@uniba.it)

**Abstract:** *Xylella fastidiosa* subsp. *pauca* ST53(*Xfp*) is responsible for Olive Quick Decline Syndrome (OQDS), which has resulted in a devastating socio-economic impact in Apulia (Southern Italy) over the last ten years. Replanting with resistant genotypes appears the most feasible and promising strategy to manage the disease to date, and the cultivars 'Leccino' and 'FS17' showed a certain level of resistance. However, no systematic and detailed information is available on neither the behaviour of the different clones of 'Leccino' towards a single or multiple infection by *Xfp*, nor the influence of certain cultural practices in modulating the resistance/susceptibility of olive trees. Moreover, a role of other endogenous factors, such as mycobiome and raw sap composition, in determining the outcome of the host/pathogen interaction cannot be excluded. The overall research plan of the PhD course aims to i) evaluate intra-cultivar variability of the cv. Leccino response towards *X. fastidiosa* infections, ii) investigate the effect of some cultural practices in modulating the resistance of different cvs towards *Xfp*, iii) verify the effect of the mycobiome and raw sap composition on the susceptibility/resistance of selected cultivars through nuclear magnetic resonance (NMR) approach, iv) investigate the diversity of wood inhabiting fungal population and their possible correlation with *Xfp* symptom severity. The results of this study, in the framework of the project ResiXO cofunded by the Regional government and CNR, could increase the chance to reduce the disease spread in Apulia and help in restoring olive cultivation in the region.

**Keywords:** mycobiome, raw sap, intra-cultivar variability, endophytes, cultural practices

## Subgroup 4 - Plant Pathology

### Preliminary data about the fungal population of Albanian apple

Amedeo Kola<sup>1\*</sup>, Annamaria Mincuzzi<sup>2</sup>, Magdalena Cara<sup>1</sup>, Marika Caputo<sup>2</sup>, Pasquale D'Ambrosio<sup>2</sup>, Jordan Merkuri<sup>3</sup>, Antonio Ippolito<sup>2</sup>

<sup>1</sup>Department of Plant Protection, Faculty of Agriculture and Environment, Agricultural University of Tirana, Albania

<sup>2</sup>Department of Soil, Plant, and Food Sciences, University of Bari Aldo Moro, Via Amendola 165/A, 70126, Bari, Italy

<sup>3</sup>Didactic and Scientific Research Center, Durres, Albania

\*Corresponding author: [akola@ubt.edu.al](mailto:akola@ubt.edu.al)

**Abstract:** In Albania the main fruit crop is represented by apple, reaching about 4,658 ha in 2021, although apple fruit is not the main production of the country. This gap is probably related to yield losses caused by pathogenic fungi. Apples are commercialized both fresh and processed, especially as juice; even if data regarding toxigenic fungi in apple fruit are almost absent, this is an important issue to human health. Mycotoxin levels are generally ruled by National and International Laws. In this study, we collected symptomatic samples of apple leaves and fruit in different regions of Albania to evaluate pathogenic fungal population. Isolates were morphologically and molecularly characterized according to fungal genus at species and morphotype level. Fungi isolated from apple leaves were *Alternaria alternata* morphotype *tenuissima* and other species within *Alternaria* genus, and *Aspergillus welwitschiae*; whereas from apple fruit *Aspergillus welwitschiae*, *Alternaria alternata* morphotype *limoniasperae*, and species belonging to *Fusarium* genus as *F. verticillioides* were isolated. Overall, according to preliminary data, *Alternaria* is the most represented genus, followed by *Fusarium* spp. Fungal distribution between apple leaves and fruit seem to be equally divided between plant organs. Abundances of species and morphotypes within *Alternaria* genus is comparable to that observed in other fruits (pomegranates, citrus, tomatoes). On fruit were also observed species belonging to *Didymellaceae*, which might be interesting for further studies.

**Keywords:** postharvest rot, wilt, alternariol, apples, *Alternaria*, *Aspergillus*, *Fusarium* spp.,

## Subgroup 5 - Weed science

### Gene flow from herbicide-tolerant hybrids to weedy sunflower

**Dragana Božić<sup>1\*</sup>, Darko Stojićević<sup>2</sup>, Sava Vrbničanin<sup>1</sup>**

<sup>1</sup>*University of Belgrade, Faculty of Agriculture, Zemun-Belgrade, Serbia*

<sup>2</sup>*Academy of Technical Vocational Studies Belgrade, Department of Applied Engineering  
Science Pozarevac, Serbia*

\*Corresponding author: [dbozic@agrif.bg.ac.rs](mailto:dbozic@agrif.bg.ac.rs)

**Abstract:** Weedy sunflower is invasive form of sunflower (*Helianthus annuus*) which widely distributed in some parts of Serbia (mainly Vojvodina province). This weed are commonly controlled by ALS-inhibiting herbicides, but wide-spread adoption of sunflower hybrids tolerant to herbicides (imazamox and tribenuron-methyl) enabled flow of genes responsible for tolerance to herbicides from that hybrids to weedy populations. The objective of this study was to check gene flow from two tolerant hybrids and estimate effect of distance between hybrid and weedy sunflower on gene flow. Gene flow was studied in the field experiments and based on DNA analysis. In the field experiments weedy sunflower were grown on different distances from tolerant hybrids. Response of progeny of weedy sunflower (collected from different distances from tolerant hybrids) to recommended rates of target herbicides were determined. As mutations responsible for sunflower tolerance to herbicides (imazamox and tribenuron-methyl) indicate gene flow, DNA analyses of selected survived plants were conducted. Although weedy sunflower progeny exposed to pollen of imazamox tolerant hybrid was survived recommended rates of herbicides, mutations were not confirmed. On the other hand gene flow from tribenuron-methyl tolerant hybrid was confirmed on molecular level. The gene responsible for tolerance was confirmed in weed sunflower samples taken from 30 and 120 m distance from the tolerant hybrid.

**Keywords:** DNA analysis, gene flow, herbicide, hybrid, weedy sunflower

## Subgroup 5 - Weed science

### Predicting *Echinochloa crus-galli* L. emergence in maize with hydrothermal modelling

Valentina Šoštarčić<sup>1\*</sup>, Roberta Masin<sup>2</sup>, Maja Šćepanović<sup>1</sup>

<sup>1</sup>University of Zagreb, Faculty of Agriculture, Svetosimunska 25, 10000 Zagreb, Croatia

<sup>2</sup>Department of Agronomy, Food, Natural Resources, Animals and Environment (DAFNAE),  
University of Padova, Viale dell'Università 16, 35020 Legnaro, Italy

\*Corresponding author: [vsostarcic@agr.hr](mailto:vsostarcic@agr.hr)

**Abstract:** Weed emergence prediction models can help determine the right time to apply herbicides and reduce herbicide input into environment. Hydrothermal models rely on soil temperature and soil moisture as key factors that promote weed emergence. A prerequisite for the use of hydrothermal models is the estimation of base temperature ( $T_b$ ) and base water potential ( $\Psi_b$ ) of the native population of weed species. In this study, the hydrothermal AlertInf developed in the Veneto region (Italy) was validated in a maize field in Croatia for the problematic weed *Echinochloa crus-galli*. Prior to validation, germination parameters ( $T_b$  and  $\Psi_b$ ) for *E. crus-galli* were estimated in laboratory experiments and then compared with Italian population built in AlertInf. The estimated  $T_b$  and  $\Psi_b$  were 0.8°C and 0.97 MPa, respectively with statistical differences only for the  $T_b$  parameter. Therefore, AlertInf was calibrated for the  $T_b$  parameter of the Croatian population. The calibrated model was used for the validation of the emergence prediction in the maize field in Croatia, Šašinoječki Lug (45°50'59.6 "N 16°09'53.9 "E) in 2019 and 2020. The overall performance of the model was evaluated using the root mean square error (RMSE) and modelling efficiency (EF). The RMSE is 1.69 and 1.38 for 2019 and 2020, respectively. In addition, EF is 0.97 and 0.98 for 2019 and 2020, respectively. The successful prediction of the calibrated model AlertInf opens the possibility to extend the model to similar geographical regions and to calibrate the model for other problematic species in the maize field in Croatia.

**Keywords:** barnyardgrass, base temperature, base water potential, integrated weed management, weeds

## Subgroup 5 - Weed science

### Secondary metabolites of *Cuscuta campestris* as bioherbicides

Tojić Teodora<sup>1\*</sup>, Marija Sarić-Krsmanović<sup>2</sup>, Tijana Đorđević<sup>2</sup>, Jelena Gajić Umiljendić<sup>2</sup>, Ljiljana Radivojević<sup>2</sup>, Dragana Božić<sup>1</sup>, Sava Vrbničanin<sup>1</sup>

<sup>1</sup>University of Belgrade, Faculty of Agriculture

<sup>2</sup>Institute of pesticides and environmental protection, Belgrade, Serbia

\*Corresponding author: [tojicteodora@gmail.com](mailto:tojicteodora@gmail.com)

**Abstract:** The harmful effects of synthetic herbicides on human health and the environment, and the increase in the number of weed species that have developed resistance to herbicides are problems that stimulated the development of new environmentally friendly approaches for weed control. A possible solution to these problems was seen in the use of the allelopathic potential of many plant species. The objective of this study was to examine the bioherbicidal potential of a secondary metabolites (phenolic compound) extracted from the stem of field dodder (*Cuscuta campestris* Yunck.) on seed germination, early seedling growth (radical and shoot length) and early plant growth tests of *Amaranthus retroflexus* and *Portulaca oleraceae*. The combined effects of the stem extract and a reduced dose of the synthetic herbicide metribuzin were also examined. The data obtained in the seed bioassay confirmed the inhibitory effect of plant stem extract on the germination and early growth of *P. oleraceae* and *A. retroflexus* seedlings at higher concentrations (0.75% and 1%). On the other hand, a synergy of field dodder stem extract and the herbicide metribuzin was revealed in plant bioassay, as their combination was found to achieve better results in the control of both weed species. Based on these obtained data, it could be concluded that *C. campestris* stem extract notably rich in phenolic acids and flavonoids could be a potential source of natural herbicide.

**Keywords:** *Cuscuta campestris*, phenolic compounds, bioherbicides, stem extracts, metribuzin.

## Subgroup 6 - Mycotoxins and food safety

### Innovative green technologies to improve the sustainability of viticulture and safety of grape products

Angelo Agnusdei, Donato Gerin, Rita Milvia Angelini De Miccolis, Stefania Pollastro\*, Francesco Faretra

Department of Soil, Plant and Food Sciences, University of Bari, via Amendola, 165/A, 70126  
Bari – Italy.

\*Corresponding author: [stefania.pollastro@uniba.it](mailto:stefania.pollastro@uniba.it)

**Abstract:** The wine sector is a driving force for the Italian economy and Apulian region is one of the leading grape growing regions in Italy. The need to adopt "green" technologies to improve the sustainability of production processes, reduce the environmental impact, rationalize the use of resources, maintaining global economic competitiveness nowadays is no longer postponable.

Phytosanitary management of vineyards is a crucial point, and it is therefore necessary to adopt new disease management strategies that lead to satisfy market demands of low environmental impact and safety and security of wine, free of the mycotoxin Ochratoxin A (OTA) and other unwanted contaminants. The objective of the research project is to integrate innovative green technologies into the grape growing process to increase the sustainability of the wine industry. The main goal will be realized through the following tasks: i) identification of new biomolecules with antimicrobial activity and biological resources with antagonistic activity; ii) development of new bio-nanocarriers to make more efficient the delivery of effective antimicrobials to the target; iii) increase knowledge on the integration of different tools to improve the effectiveness of crop protection strategies and prevent new phytosanitary emergencies; iv) improve knowledge on biosynthesis and biological role of Ochratoxin A; v) improve the quantitative detection of OTA producing *Aspergillus carbonarius* in the lab and even directly in the field through portable devices; vi) validation of disease forecasting models for vineyards in southern Italy.

**Keywords:** Ochratoxin A, *Aspergillus carbonarius*, plant protection products, nanocarriers, wine



## Subgroup 6 - Mycotoxins and food safety

### Cyclopiazonic acid production in *Penicillium* spp. causing postharvest pome fruit decay

Aleksandra Žebeljan\*, Nataša Duduk, Nina Vučković, Ivana Vico

University of Belgrade - Faculty of Agriculture, Nemanjina 6, Belgrade, Serbia

\*Corresponding author: [zheki24@gmail.com](mailto:zheki24@gmail.com)

**Abstract:** *Penicillium* spp, are diverse and cosmopolite fungi which cause blue mold, one of the most economically important postharvest diseases of pome fruit, worldwide. In addition to causing fruit decay, *Penicillium* spp. are known to produce a range of mycotoxins including patulin, ochratoxin A, citrinin, penitrem A, roquefortine, citreoviridin, and cyclopiazonic acid. Cyclopiazonic acid is also an extrolite useful in the identification of a *Penicillium* isolate. In Serbia, *Penicillium* spp. responsible for postharvest decay of pome fruit (apple, pear, quince, and medlar) are *P. expansum*, *P. crustosum*, and *P. solitum*. The production of cyclopiazonic acid in a total of 26 *Penicillium* isolates from pome fruit belonging to three species: *P. expansum*, *P. crustosum*, and *P. solitum* was examined using Ehrlich's test (Lund, 1995). Filter paper emersed in Ehrlich reagent was placed on top of the mycelial side of agar plugs and after incubation, reaction was recorded based on the appearance of color. Amongst *P. expansum*, 12 out of 18 isolates formed yellow rings and six isolates formed faint violet rings. All seven *P. crustosum* isolates formed faint yellow to yellow rings and *P. solitum* isolate also formed a yellow ring. Observed reactions indicated that six *P. expansum* isolates produced cyclopiazonic acid, while the other 12 *P. expansum* isolates, and all *P. crustosum* and *P. solitum* isolate produced other alkaloids.

**Keywords:** Ehrlich's test, mycotoxins, *Penicillium expansum*, *Penicillium crustosum*, *Penicillium solitum*



## Subgroup 6 - Mycotoxins and food safety

### Occurrence of ochratoxin A in Blatina wine from the Mostar vineyards

Ana Sesar<sup>1</sup>, Anita Lalić<sup>2\*</sup>, Kristina Batinić<sup>1</sup>

<sup>1</sup>Federal Agro-Mediterranean Institute, Mostar, 88 000 Mostar, Bosnia and Herzegovina

<sup>2</sup>Faculty of Agriculture and Food Technology, 88 000 Mostar, Bosnia and Herzegovina

\*Corresponding author: [anita.juric@aptf.sum.ba](mailto:anita.juric@aptf.sum.ba)

**Abstract:** Blatina is an autochthonous and most represented red wine variety in Herzegovina. Wine is the second source of daily micotoxins intake for humans with domination of ochratoxin A (OTA) contamination of grapes. The aim of this paper was to determine occurrence of OTA in Blatina. Representative sampling was achieved by selecting twenty quality and premium wines from Mostar vineyard produced in 2018 and 2019. Quantification of OTA was performed by high performance liquid chromatography with a fluorescent detector (HPLC-FLD). OTA values ranged from 0.205 to 0.933 µg/L. It can be concluded that the presence of ochratoxin A in Blatina wine is within acceptable limits of Regulation in the European Union (EU). The level of ochratoxin A in wine depends on the type of grape, wine region, agricultural practice, weather conditions and the process of wine production. In order to protect consumer health, it is necessary to monitor its values in wines on the market.

**Keywords:** red wine, ochratoxin A, HPLC-FLD

## Subgroup 6 - Mycotoxins and food safety

### Biological control of aflatoxins using non-toxigenic strains of *Aspergillus flavus* on maize

Maria Varveri, Mike Kaminariis, Spyridoula Dimitropoulou, Dimitrios Tsitsigiannis\*

Department of Crop Science, Laboratory of Plant Pathology

Agricultural University of Athens, Greece

\*Corresponding author: [dimtsi@aua.gr](mailto:dimtsi@aua.gr)

**Abstract:** Mycotoxins are one of the major threats to food and feed safety and quality worldwide. More specifically, aflatoxin AFB1 and its metabolite AFM1, have been classified by the International Agency for Research on Cancer (IARC) among the most carcinogenic compounds for humans. Due to the inability of chemical methods to control aflatoxin levels on maize, the use of non-toxigenic strains of *Aspergillus flavus* has been characterized by numerous studies as the most effective control strategy. The purpose of the present study was to evaluate 15 non-toxigenic strains in terms of their ability to reduce aflatoxin production *in situ*, on artificially infected corn seeds with a highly toxigenic *A. flavus* strain from the collection of the Laboratory of Phytopathology, AUA. Previous experiments have indicated the high effectiveness of these specific non-toxigenic isolates in inhibiting the biosynthesis of aflatoxins on pistachios, reaching reduction rates of aflatoxin levels between 80-90% both in laboratory experiments and in field experiments. In the context of biological management, the most effective of the non-toxigenic isolates in corn kernels will be further applied to maize cultivation, under field conditions in order to further evaluate their effectiveness. The results of the above experimental studies will be presented.

**Keywords:** *Aspergillus flavus*, aflatoxins, biological control, non-toxigenic strains, maize

## Subgroup 7 - Bio-diversity and bio-indicators in sustainable agriculture

### Neglected and Underutilized Crop Species, an alternative to improve food supply and nutrition security by protecting biodiversity in Korça region, Albania

Robert Damo, Pirro Icka\*

Department of Agronomy, University Fan Noli Korça, Albania

\*Corresponding author [pirroicka@yahoo.com](mailto:pirroicka@yahoo.com)

**Abstract:** Agrobiodiversity as part of humanity cultural heritage is essential for a sustainable agriculture production, food, and nutritional security, of which neglected and underutilized species are key elements. Many of these species are well adapted to specific regions, to various environmental stresses and have good *resistance* to common *diseases and pests*. They have a high potential for providing food, and nutritional security to support local food system transformation and contribute to the diversity and stability of agro-ecosystems and the diversification of agriculture. Korça region characterized by diverse agro-ecology and it considered as one of the richest crop genetic resource centers in Albania. The cultivation and use of minor crops is declining from time to time and nowadays have limited attention on the diversity and conservation of those genetic resources. As a result, some of them are endangered, and to be lost in near future. The aim of this paper is to indicate several crop species that are neglected or underutilized in Korça region, which can be included and used as future smart foods, and may contribute on diversification of food and cultivated crops.

**Keywords:** neglected crop, underutilized crop, biodiversity, future smart foods.

## Subgroup 7 - Bio-diversity and bio-indicators in sustainable agriculture

### Nematodes as bioindicators of soil ecosystem status in the intercropping system of woody species and agricultural crops

Josipa Puškarić<sup>1\*</sup>, Vladimir Ivezić<sup>1</sup>, Brigita Popović<sup>1</sup>, Dinka Grubišić<sup>2</sup>, Milan Radivojević<sup>3</sup>, Nikola Grujić<sup>3</sup>, Mirjana Brmež<sup>1</sup>

<sup>1</sup>Josip Juraj Strossmayer University of Osijek, Faculty of Agrobiotechnical sciences Osijek, Osijek, Croatia, <sup>2</sup>University of Zagreb, Faculty of Agriculture, Zagreb, Croatia, <sup>3</sup>University of Belgrade, Faculty of Agriculture, Belgrade, Serbia

\*Corresponding author: [josipa.puskaric@fazos.hr](mailto:josipa.puskaric@fazos.hr)

**Abstract:** The aim of this study was to determine the impact of intercropping of permanent plantation and agricultural crops on biodiversity, community structure and soil disturbance indices and ecological diversity indices as well as on fertility parameters. For the purpose of the research, experiments were conducted on three different ecosystems, the intercropping ecosystem of agricultural crops and walnut (K+O), the walnut ecosystem (O) and the agricultural crops ecosystem (K), on two sites (Ivankovo and Đakovo), from 2017 to 2019. The ecosystem K+O has shown a positive impact on total and average diversity of genera, especially on the diversity of useful omnivores, which indicates the stability of the ecosystem. Bacterivores were significantly more numerous in the ecosystem K, and fungivore in the ecosystem O. The ecosystem K+O indicates the decomposition of organic matter by both bacterial and fungal pathways. The values of the MI and MI 2-5 indicates the greatest disturbances in the ecosystem K. The CI values in the ecosystem O indicate a higher proportion of fungi and fungivores that participate in the decomposition of organic matter. BI and EI values in all ecosystems indicate that soil disturbance is present, but the soil food network is not depleted. SI values indicate the developed soil food web and a good share of higher trophic groups in ecosystems K+O and O. Metabolic footprints of nematodes, high activity of bacteria and fungi, higher amounts of organic matter and soil moisture also indicate a positive impact of the ecosystem K+O, when compared to the ecosystem K.

**Keywords:** nematodes, biodiversity, indices, intercropping, walnut



## Program

### Program of the conference

#### Color index:

New day	Section name
Hall	Breaks

#### SG index:

SG 1 - Diagnosis in plant health and IPM	SG 2 - Sustainable use of pesticides
SG 3- Subgroup 3 - Plant feeders	SG 4 - Plant Pathology
SG 5 - Weed science	SG 6 - Mycotoxins and food safety
SG 7 - Bio-diversity and bio-indicators in sustainable agriculture	

#### Monday, September 5<sup>th</sup> 2022

15:00-20:00 Participants arrival

#### Tuesday, September 6<sup>th</sup> 2022

Faculty of Agriculture, University of Novi Sad  
Trg Dositeja Obradovica 8, Novi Sad

9:00-10:00 Registration of participants

10:00-10:15

#### Opening Session

Amphitheatre, ground floor

**Prof. Aleksandra Konjević PhD**, local project coordinator, president of the organising committee

**Prof. Branko Ćupina PhD**, Vice Dean for Science and International Relations, Faculty of Agriculture, University of Novi Sad

10:15-13:15

#### "Presentation of WPs project results "

Amphitheatre, ground floor

10:15-10:35 **Prof. Renata Bažok PhD**, project leader: Project HarISA, short overview

10:35-10:50 **Prof. Vili Harizanova PhD**, leader of WP2: General rules

10:50-11:20 **Prof. Aleksandra Ignjatović-Ćupina PhD**, **Prof. Eustachio Tarasco PhD**, leaders of WP3: Curriculum of the Joint PhD study Program

11:20-11:30 **Discussion**

11:30-12:00 Coffee break

12:00-12:15 **Prof. Dimitrios Tsitsigiannis PhD**, **Prof. Magdalena Cara PhD**, leaders of WP4: Equipment and contracts

12:15-12:30 **Prof. Renata Bažok PhD**, **Prof. Ilir Nicko PhD**, leaders of WP5: Mobility of students and professors

12:30-12:45 **Prof. Marija Grgić PhD**, leader of WP6: Quality Control Board report

12:45-13:00 **Prof. Renata Bažok PhD**, **Prof. Jurica Primorac PhD**, leaders of WP7: Dissemination of project activities and results

13:00-13:15 **Discussion – How to proceed?**

13:15-14:45 Lunch break



## Program

Color index:		SG index:	
New day	Section name	SG 1 - Diagnosis in plant health and IPM	SG 2 - Sustainable use of pesticides
		SG 3- Subgroup 3 - Plant feeders	SG 4 - Plant Pathology
Hall	Breaks	SG 5 - Weed science	SG 6 - Mycotoxins and food safety
		SG 7 - Bio-diversity and bio-indicators in sustainable agriculture	

Wednesday, September 7 <sup>th</sup> 2022		
Faculty of Agriculture, University of Novi Sad Trg Dositeja Obradovica 8, Novi Sad		
9:30-11:45	"Lectures" – part 2 - Parallel sessions	
	P1, ground floor, SG3 Chaired by: <b>Ivana Majić, Nedžad Karić</b>	Amphitheatre, ground floor, SG4 Chaired by: <b>Sotiris Tjamos, Nataša Duduk</b>
9:30-9:45	<b>Renata Bažok:</b> Effects of spiropidion on survival and parasitism of <i>Aphidius collemanni</i> Viereck (Hymenoptera: Braconidae)	<b>Rita Milvia De Miccolis Angelini:</b> Multi-omics approaches to explore phenotypic and genetic diversity in the brown rot fungal pathogens <i>Monilinia fructicola</i> , <i>Monilinia laxa</i> and <i>Monilinia fructigena</i>
9:45-10:00	<b>Martina Kadoić Balaško:</b> Electrical penetration graph - Promising tool for measuring feeding behaviour of <i>Bemisia tabaci</i>	<b>Tatjana Dudaš:</b> Biocontrol activity of epiphytic yeasts against <i>Penicillium expansum</i> on apple fruits
10:00-10:15	<b>Ilaria Laterza:</b> Impacts of landscape composition on the occurrence and the control of <i>Halyomorpha halys</i> in newly invaded areas	<b>Almira Konjić:</b> Overview of the use of GWAS to identify loci for resistance to apple scab caused by <i>Venturia inaequalis</i>
10:15-10:30	<b>Aleksandar Ivezić:</b> Identification of Trichogramma egg parasitoids of the European corn borer <i>Ostrinia nubilalis</i> (Hübner, 1796) in Vojvodina (Serbia)	<b>Anastasia Venieraki:</b> Antagonistic activity of endophytic <i>Bacillus</i> strains against phytopathogenic fungi
10:30-10:45	<b>Sofia Dervisoglou:</b> <i>Tuta absoluta</i> egg density and distribution on tomato plants affect the efficacy of mirid predators	<b>Eirini Poulaki:</b> The volatile organic compounds of the biocontrol agent <i>Bacillus velezensis</i> K165 activate plant defenses against <i>Verticillium dahliae</i>
10:45-11:00	<b>Ivana Lalićević:</b> Aphid flight activity in wheat	<b>Christos Tsoukas:</b> Grapevine rootstock and scion canes as an important inoculum source of Petri disease associated pathogens
11:00-11:15	<b>Marija Milošević:</b> Natural enemies of <i>Parthenolecanium corni</i> (Bouché) (Hemiptera: Coccidae) on grapevine in Neštin, Serbia	<b>Nina Vučković:</b> Botryosphaeriaceae as postharvest pathogens of apple fruit

## Program

11:15-11:30	Mihaela Kavran: Some important invasive species in agricultural and urban environments of Serbia	Discussion
11:30-11:45	Short discussion	
11:45-12:15	Coffee break	
12:15-13:30	"Lectures" – part 2	
	P1, ground floor, SG5 Chaired by: Maja Šćepanović, Mirha Đikić	Amphitheatre, ground floor, SG4, SG6 Chaired by: Dimitrios Tsitsigiannis, Magdalena Cara
12:15-12:30	Dragana Božić: Gene flow from herbicide-tolerant hybrids to weedy sunflower	Anastasia Papageorgiou: Endophytic bacteria from olive drupes as biological control and induced resistance agents of <i>Colletotrichum acutatum</i>
12:30-12:45	Valentina Šoštarčić: Predicting <i>Echinochloa crus-galli</i> L. emergence in maize with hydrothermal modelling	Angelo Agnusdei: Innovative green technologies to improve the sustainability of viticulture and safety of grape products
12:45-13:00	Teodora Tojić: Secondary metabolites of <i>Cuscuta campestris</i> as bioherbicides	Discussion
13:00-13:15	Short discussion	
13:15-14:45	Lunch	
14:45-16:30	Poster session	
	Amphitheatre, hall, ground floor Chaired by: Mirjana Brmež, Nedeljko Latinović	Meeting hall, dean's office
14:45-15:15	Online posters presentation and discussion	WP2 and PMB meeting
15:15-16:30	Poster session on site	

## Program

Color index:		SG index:	
New day	Section name	SG 1 - Diagnosis in plant health and IPM	SG 2 - Sustainable use of pesticides
		SG 3- Subgroup 3 - Plant feeders	SG 4 - Plant Pathology
Hall	Breaks	SG 5 - Weed science	SG 6 - Mycotoxins and food safety
		SG 7 - Bio-diversity and bio-indicators in sustainable agriculture	

### Thursday, September 8<sup>th</sup> 2022

Faculty of Agriculture, University of Novi Sad  
Trg Dositeja Obradovica 8, Novi Sad

<b>9:00-11:30</b>	<b>Students' mobility presentations and Final discussion</b>	
	Amphitheatre, ground floor <b>Chaired by: Karolina Vrandečić, Marta Loc</b>	P1, ground floor
9:00-11:00	<b>Round table:</b> Student's experience and feelings about the mobilities	<b>PMB meeting</b>
11:00-11:30	<b>Discussion about future activities and conclusions</b>	
11:30-12:00	Coffee break	
<b>12:00</b>	<b>Field trip</b>	

### Friday, September 9<sup>th</sup> 2022

	<b>Departures</b>
--	-------------------

***This book is made with the support of:***

- ✓ Erasmus+ KA2 Program of the European Union



Co-funded by the  
**Erasmus+ Programme**  
of the European Union

- ✓ Republic of Serbia, Autonomous Province of Vojvodina, Provincial Secretariat for Finance



***Conference organizers:***

- ✓ University of Novi Sad, Faculty of Agriculture, Serbia



- ✓ University of Zagreb, Faculty of Agriculture, Croatia

