


# 2302272 – Org Chem II – Part IV

## Lecture 1

### Protein-1

---



*Instructor: Asst. Prof. Dr. Tanatorn Khotavivattana*

*E-mail: [tanatorn.k@chula.ac.th](mailto:tanatorn.k@chula.ac.th)*

#### **Recommended Textbook:**

Chapter 24 in *Organic Chemistry*, 8<sup>th</sup> Edition, L. G. Wade, Jr., 2010, Prentice Hall (Pearson Education)

# Key concepts:





- 1) โพรตีนคืออะไร? เจอได้ที่ไหนบ้าง? ทำหน้าที่อะไรได้บ้าง?
- 2) ต้องมีโครงสร้างแบบไหนถึงจะเรียกว่าเป็นโปรตีน?  
กรดอะมิโนเกี่ยวข้องกับโปรตีน?

# Protein

Proteins are the **most abundant** organic molecules in **animals**, playing important roles in all aspects of cell structure and function




Class of Protein	Example	Function of Example
structural proteins	collagen, keratin	strengthen tendons, skin, hair, nails
enzymes	DNA polymerase	replicates and repairs DNA
transport proteins	hemoglobin	transports O <sub>2</sub> to the cells
contractile proteins	actin, myosin	cause contraction of muscles
protective proteins	antibodies	complex with foreign proteins
hormones	insulin	regulates glucose metabolism
toxins	snake venoms	incapacitate prey



# Protein – General structure

- Proteins are **biopolymers** of  **$\alpha$ -amino acids**, so named because the amino group is bonded to the  $\alpha$  carbon atom, next to the carbonyl group




- The amino acid subunits are joined by **amide linkages** called **peptide bonds**

# Key concepts:


3) กรดอะมิโนแต่ละชนิดแตกต่างกันอย่างไร?

# $\alpha$ -Amino Acids

- The simplest  $\alpha$ -amino acid is **aminoacetic acid**, called **glycine**. Other common amino acids have **side chains** (symbolised by **R**) substituted on the  **$\alpha$  carbon**; for example, **alanine** is the amino acid with a **methyl side chain**



- Except for glycine, the  $\alpha$ -amino acids are all **chiral** and the chirality centre is at the asymmetric  $\alpha$  carbon atom
- Nearly all the naturally occurring amino acids are found to have the **(S) configuration**



Only this isomer occurs in proteins.

# Standard $\alpha$ -Amino Acids

20 **standard** amino acids, grouped according to the chemical properties of side chains

Name	Symbol	Abbreviation	Structure	Functional Group in Side Chain	Isoelectric Point
side chain is nonpolar, H or alkyl					
glycine	G	Gly		none	6.0
alanine	A	Ala		alkyl group	6.0
*valine	V	Val		alkyl group	6.0
*leucine	L	Leu		alkyl group	6.0
*isoleucine	I	Ile		alkyl group	6.0
*phenylalanine	F	Phe		aromatic group	5.5
proline	P	Pro		rigid cyclic structure	6.3
*essential amino acid					

Name	Symbol	Abbreviation	Structure	Functional Group in Side Chain	Isoelectric Point
<b>side chain contains an —OH</b>					
serine	S	Ser	$\begin{array}{c} \text{HOOC} \\   \\ \text{H}_2\text{N} \text{---} \text{C} \text{---} \text{H} \\   \\ \text{H} \end{array}$	hydroxyl group	5.7
*threonine	T	Thr	$\begin{array}{c} \text{HOOC} \\   \\ \text{H}_2\text{N} \text{---} \text{C} \text{---} \text{H} \\   \\ \text{H} \end{array}$	hydroxyl group	5.6
tyrosine	Y	Tyr	$\begin{array}{c} \text{HOOC} \\   \\ \text{H}_2\text{N} \text{---} \text{C} \text{---} \text{H} \\   \\ \text{H} \end{array}$	phenolic—OH group	5.7
<b>side chain contains sulfur</b>					
cysteine	C	Cys	$\begin{array}{c} \text{HOOC} \\   \\ \text{H}_2\text{N} \text{---} \text{C} \text{---} \text{H} \\   \\ \text{H} \end{array}$	thiol	5.0
*methionine	M	Met	$\begin{array}{c} \text{HOOC} \\   \\ \text{H}_2\text{N} \text{---} \text{C} \text{---} \text{H} \\   \\ \text{H} \end{array}$	sulfide	5.7
<b>side chain is acidic</b>					
aspartic acid	D	Asp	$\begin{array}{c} \text{HOOC} \\   \\ \text{H}_2\text{N} \text{---} \text{C} \text{---} \text{H} \\   \\ \text{H} \end{array}$	carboxylic acid	2.8
glutamic acid	E	Glu	$\begin{array}{c} \text{HOOC} \\   \\ \text{H}_2\text{N} \text{---} \text{C} \text{---} \text{H} \\   \\ \text{H} \end{array}$	carboxylic acid	3.2



Name	Symbol	Abbreviation	Structure	Functional Group in Side Chain	Isoelectric Point
<b>side chain is basic</b>					
*lysine	K	Lys	$\begin{array}{c} \text{HOOC} \\   \\ \text{H}_2\text{N} \text{---} \text{C} \text{---} \text{H} \\   \\ \text{H} \end{array}$	amino group	9.7
*arginine	R	Arg	$\begin{array}{c} \text{HOOC} \\   \\ \text{H}_2\text{N} \text{---} \text{C} \text{---} \text{H} \\   \\ \text{H} \end{array}$	guanidino group	10.8
*histidine	H	His	$\begin{array}{c} \text{HOOC} \\   \\ \text{H}_2\text{N} \text{---} \text{C} \text{---} \text{H} \\   \\ \text{H} \end{array}$	imidazole ring	7.6
<b>side chain contains nonbasic nitrogen</b>					
asparagine	N	Asn	$\begin{array}{c} \text{HOOC} \\   \\ \text{H}_2\text{N} \text{---} \text{C} \text{---} \text{H} \\   \\ \text{H} \end{array}$	amide	5.4
glutamine	Q	Gln	$\begin{array}{c} \text{HOOC} \\   \\ \text{H}_2\text{N} \text{---} \text{C} \text{---} \text{H} \\   \\ \text{H} \end{array}$	amide	5.7
*tryptophan	W	Trp	$\begin{array}{c} \text{HOOC} \\   \\ \text{H}_2\text{N} \text{---} \text{C} \text{---} \text{H} \\   \\ \text{H} \end{array}$	indole	5.9

# Key concepts:

- 4) กรดอะมิโนแต่ละชนิดมีความเป็นกรด/เบส ต่างกันอย่างไร?
- 5) Isoelectric point คืออะไร?  
เกี่ยวข้องกับอย่างไรกับ Electrophoresis?




## $\alpha$ -Amino Acids – Acid-Base Properties

- Although we commonly write amino acids with an intact carboxyl  $-\text{COOH}$  group and amino  $-\text{NH}_2$  group, their actual structure is **ionic** and **depends on the pH**
  
- This structure is called a **zwitterion**, giving them some **unusual properties**
  - 1) More **soluble in water** than they are in common organic solvents
  - 2) **High melting points**, generally over  $200\text{ }^\circ\text{C}$
  - 3) Much **larger dipole moments** than simple amines or simple acids

## $\alpha$ -Amino Acids – Acid-Base Properties

- This structure is called a **zwitterion**, giving them some **unusual properties**


**4) Less acidic than most carboxylic acids and less basic than most amines**



- Because amino acids contain both acidic ( $-\text{NH}_3^+$ ) and basic groups ( $-\text{CO}_2^-$ ), they are **amphoteric** (having both acidic and basic properties).

# $\alpha$ -Amino Acids – Acid-Base Properties

- The predominant form of the amino acid depends on the **pH** of the solution





- In an **acidic** solution, the  $\text{---COO}^-$  group is **protonated** to a free  $\text{---COOH}$  group, and the molecule has an overall **positive charge**
- As the pH is raised, the  $\text{---COOH}$  loses its proton and becomes **zwitterion** at about pH 2. This point is called  **$pK_{a1}$ , the first acid-dissociation constant**
- As the pH is raised further, the  $\text{---NH}_3^+$  group loses its proton at about pH 9 or 10. This point is called  **$pK_{a2}$ , the second acid-dissociation constant**. Above this pH, the molecule has an overall **negative charge**

# $\alpha$ -Amino Acids – Acid-Base Properties

- Titration curve for glycine;


**Isoelectric point (pI)** is the pH where the amino acid exists in the zwitterionic form



# $\alpha$ -Amino Acids – Acid-Base Properties

- Isoelectric pH depends on the amino acid structure

**Neutral amino acids:** (5.0 to 6.3); **alanine (6.0)**



**Acidic amino acids:** **aspartic acid (2.8)**, glutamic acid (3.2)

**Basic amino acids:** **lysine (9.7)**, arginine (10.8), histidine (7.6)

# Electrophoresis


separate mixtures of amino acids

12


An amino acid mixture is placed in the centre of a gel. Two electrodes are placed at the edges, and a potential is applied across the electrodes

**Positively charged** amino acids are attracted to the **negative electrode**, and **negatively charged amino acids** are attracted to the **positive electrode**. An amino acid at its **isoelectric** point has no net charge, so it **does not move**


*Beginning*




*End*




*Structure at pH 6*



alanine (charge )



lysine (charge )



aspartic acid (charge )




# Key concepts:

6) ถ้าอยากสร้างกรดอะมิโนขึ้นมาเอง ไม่ได้สกัดหรือแยกมาจากธรรมชาติ จะทำได้อย่างไร?


# $\alpha$ -Amino Acids – Synthesis

1) **Reductive Amination**; formation of amino acids from appropriate  **$\alpha$ -ketoacid**




# $\alpha$ -Amino Acids – Synthesis

## 2) $S_N2$ Amination of $\alpha$ -haloacid



# $\alpha$ -Amino Acids – Synthesis

3) **The Strecker Synthesis**; formation of amino acids from appropriate **aldehydes**




# Key concepts:

7) เราสามารถปรับเปลี่ยนโครงสร้างของกรดอะมิโนได้อย่างไรบ้าง?

# $\alpha$ -Amino Acids – Reactions

## 1) Esterification of the Carboxyl Group


- Like other carboxylic acids, amino acids are **esterified** by treatment with a large excess of an **alcohol** and an **acidic catalyst**; Under acidic conditions, the amino group is present in its protonated form, so it does not interfere with esterification



# $\alpha$ -Amino Acids – Reactions

## 1) Esterification of the Carboxyl Group

- Esters of amino acids are often used as **protected derivatives** to prevent the carboxyl group from reacting in some undesired manner
- **Aqueous acid** hydrolyses the ester and **regenerates the free amino acid**



# $\alpha$ -Amino Acids – Reactions

## 2) Acylation of the Amino Group – Formation of Amides

- **Acylating agent** (acid chloride/anhydride) converts the amino group to an **amide**






# $\alpha$ -Amino Acids – Reactions

## 2) Acylation of the Amino Group – Formation of Amides

- **Benzyl chloroformate** acylates the amino group to give a benzyloxycarbonyl derivative, often used as a **protecting group in peptide synthesis**




# $\alpha$ -Amino Acids – Reactions

## 3) Reaction with Ninhydrin

- Ninhydrin is a common reagent for **visualizing** spots or bands of amino acids that have been separated by chromatography or electrophoresis

*Reaction of an amino acid with ninhydrin*



- When ninhydrin reacts with an amino acid, one of the products is a deep violet, resonance-stabilised anion called **Ruhemann's purple**
- Ninhydrin produces this same purple dye regardless of the structure of the original amino acid
- The side chain of the amino acid is lost as an **aldehyde**

