

Time left 0:04:28

1. **About smoking, which statement is incorrect?**

- a. Most of those killed by tobacco are not particularly heavy smokers and most started as teenagers
- b. Number of carcinogenic substances associated with smoking is about 70
- c. Approximately 50 percent of smokers die prematurely from their smoking, on average 8 years earlier than non-smokers
- d. There is evidence that smoking can cause about 40 different diseases
- e. Number of harmful substances associated with smoking is about 7240

[Clear my choice](#)

**Collected by:
Maria Maghaireh**

[Next page](#)

Time left 0:04:28

1. **About smoking, which statement is incorrect?**

- a. Most of those killed by tobacco are not particularly heavy smokers and most started as teenagers
- b. Number of carcinogenic substances associated with smoking is about 70
- c. Approximately 50 percent of smokers die prematurely from their smoking, on average 8 years earlier than non-smokers
- d. There is evidence that smoking can cause about 40 different diseases
- e. Number of harmful substances associated with smoking is about 7240

[Clear my choice](#)

[Next page](#)

Confirmed case definition is made based on ..

- a. Diagnostic lab. Workup.
- b. Contact with other probable cases.
- c. Contacts with other cases.
- d. Clinical Symptoms.

[Clear my choice](#)

Time le

Next

Time left 0:10:45

Which of the following is associated with bioterrorism?

- a. Re-emerging disease.
- b. Zoonotic jump.
- c. Deliberately emerging diseases.
- d. Accidentally emerging.

Clear my choice

Next page

Question **43**

Not yet
answered

Marked out of
1.0

Prevalence is the actual number of cases alive and dead

Select one:

True

False

Next page

طب مجت

es / طب مجتمع (1) ش 1 / General / Med 410 Midterm

Time left 0:13:52

Concerning grand challenges for mental disorders, strengthen the mental health component in the training of all healthcare personnel is related to the goal of improving treatments and expand access to care

Select one:

True

False

Next page

Search



Which of the following infectious diseases may have the largest prevalence rate?

- a. Flu.
- b. Common cold.
- c. HIV.
- d. COVID. 19.

5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
0	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46

Time left 0:15:20

about the impact of non-communicable diseases on global burden of diseases, which statement is incorrect?

- a. The number of patients with diabetes in the world is expected to increase from 194 Million in 2003 to 330 in 2030 with three in four living in developing countries
- b. In some countries of the Middle East, one in four deaths in adults aged between 35 and 64 years is attributable to diabetes
- c. In developed countries most people with diabetes are above the age of retirement, whereas in developing countries those most frequently affected are aged between 25 and 46
- d. By 2020, it is predicted that these diseases will be causing seven out of every 10 deaths in developing countries

Clear my choice

Question 33
Not yet answered
Marked out of 1.0

Time left 0:13:38

About mental health, which statement is incorrect?

- a. stress is a response to an external cause
- b. Anxiety is a person's specific reaction to stress; its origin is internal
- c. People with an anxiety disorder are three to five times more likely to go to the doctor and six times more likely to be hospitalized for psychiatric disorders than those who do not suffer from anxiety disorders
- d. It's estimated that job stress-costs U.S. industry more than \$300 billion a year in absenteeism, turnover, diminished productivity, and medical, legal, and insurance costs
- e. Anxiety is the body's reaction to a threat, whereas stress is the body's reaction to the anxiety

Clear my choice

Quiz navigation

1	2
10	11
19	20
28	29
37	38
46	47

Finish attempt



What will happen for an epidemic with reproductive rate of 1.5 ..

- a. Infection will widely spread.
- b. Infection will last for long time.
- c. Infection will not widely spread.

Clear my choice

5	6	7	8	9	10	11	12	13	14	15	16	17	18
10	31	32	33	34	35	36	37	38	39	40	41	42	43

Time left 0:16:40

Question 29

Not yet answered

Marked out of 1.0

A man with a standard life expectancy of 75 years died at age of 60. When he was at age of 40 he had amputation from diabetic foot therefore he spent the rest of his life in a disability state with a weight factor of 0.5, The DALYs for this person is:

- a. 10 years
- b. 20 years
- c. 25 years
- d. 15 years

[Clear my choice](#)

[Next page](#)

The uses for Public Health Surveillance include the followings except

- a. Estimate magnitude and scope of health problems
- b. Identify patients and their contacts for treatment and intervention
- c. Develop hypotheses and stimulate research
- d. Assess effectiveness of programs and control measures
- e. Detect the suitable treatment for patients

[Clear my choice](#)

5	6	7	8	9	10	11	12	13	14	15	16	17	18	19
20	31	32	33	34	35	36	37	38	39	40	41	42	43	44

Question **27**
Not yet
answered
Marked out of
1.0

Multiple paid media (TV, radio, print, etc.), public relations, special events and promotions are related to efforts to maximize the success of any program to reduce tobacco use

- a. Community-Based Programs
- b. Related Policy Efforts
- c. Public Education Efforts
- d. Monitoring and Evaluation

Clear my choice

Next page



Time left 0:18:32

Question 26

Not yet answered

Marked out of 1.0

According to CDC, functions of epidemiology include the followings except:

- a. Offer Interventions
- b. Public Health Surveillance
- c. Generate epidemic hypothesis
- d. Communication
- e. Evaluation of disease control programs

[Clear my choice](#)

[Next page](#)



طب مجتمع

My courses / طب مجتمع (1) ش1 / General / Med 410 Midterm


Time left 0:

Human made factor that is strongly associated with the rise of wide range pandemic like COVID-19 ..

- a. Globalization and traveling behaviors.
- b. Dam building.
- c. Changes in food industry.
- d. Urbanization.

[Clear my choice](#)

Next page



Med 410 Midterm (page 28 of 50) x

moodle.yu.edu.jo/exam/mod/quiz/attempt.php?attempt=47202&cmid=11515&page=27

YouTube Maps

Exam System English (en)

طب مجتمع ()

My courses / طب مجتمع / General / Med 410 Midterm


Time left 0:23:09

Which of the following is associated with bioterrorism?

- a. Accidentally emerging.
- b. Zoonotic jump.
- c. Deliberately emerging diseases.
- d. Re-emerging disease.

Clear my choice

Next page



طب مجتمع

courses / طب مجتمع (1) ش1 / General / Med 410 Midterm

Time left 0:22:14

About mental health, which statement is incorrect?

- a. Anxiety is the body's reaction to a threat, whereas stress is the body's reaction to the anxiety
- b. stress is a response to an external cause
- c. People with an anxiety disorder are three to five times more likely to go to the doctor and six times more likely to be hospitalized for psychiatric disorders than those who do not suffer from anxiety disorders
- d. Anxiety is a person's specific reaction to stress; its origin is internal
- e. It's estimated that job stress costs U.S. industry more than \$300 billion a year in absenteeism, turnover, diminished productivity, and medical, legal, and insurance costs

pe here to search



طب مجتمع (1) ش

Dashboard / My courses / طب مجتمع (1) ش / General / Med 410 Midterm

Time left 0:25

Question 25
Correctly answered
Marked out of

Which of the followings is not among the activities of descriptive epidemiology:

- a. To plan, provide, and evaluate health services
- b. Study the course of disease within a geographical region
- c. Answer why and how an epidemic is existing
- d. Generate hypotheses
- e. Assess who is affected by the health problem

Clear my choice

Next page

TOSHIBA



Question **24**

Not yet answered

Marked out of 1.0

Which of the followings is not among the activities of descriptive epidemiology:

- a. To plan, provide, and evaluate health services
- b. Answer why and how an epidemic is existing
- c. Study the course of disease within a geographical region
- d. Assess who is affected by the health problem
- e. Generate hypotheses

[Clear my choice](#)

[Next page](#)



Time left 0:24:45

An epidemic curve with multiple peaks overtime is usually the result of what type of exposure?


- a. Continuous common source.
- b. Point source.
- c. Intermittent common source.
- d. Person to person transmission.

[Clear my choice](#)

Next page

exam/mod/quiz/attempt.php?attempt=47185&cmid=11515&page=10

here to search



طب مجتمع

...y courses / طب مجتمع (1) / General / Med 410 Midterm

Time left 0:26

Which of the following is associated with the development of new variant of the same agent?

- a. Deliberately emerging diseases.
- b. Accidentally emerging.
- c. Zoonotic jump.
- d. Re-emerging disease.

[Clear my choice](#)

Next page



TOSHIBA




courses / 1 (1)

An epidemic curve with multiple peaks overtime is usually the result of what type of exposure?

- a. Person to person transmission.
- b. Point source.
- c. Continuous common source.
- d. Intermittent common source.

[Clear my choice](#)



Time left 0:24:30

Question **33**

Not yet
answered

Marked out of
1.0

Which of the following intervals or time period is important for infectious agent to reproduce and replicate in the host body?

- a. Latent period.
- b. Infectious period.
- c. Incubation period.
- d. Serial interval.

[Clear my choice](#)



[Next page](#)

Time left (

About the impact of non-communicable diseases on global burden of diseases, which statement is incorrect?

- a. In developed countries most people with diabetes are above the age of retirement, whereas in developing countries those most frequently affected are aged between 35 and 64
- b. In some countries of the Middle East, one in four deaths in adults aged between 35 and 64 years is attributable to diabetes
- c. By 2020, it is predicted that these diseases will be causing seven out of every 10 deaths in developing countries
- d. The number of patients with diabetes in the world is expected to increase from 194 Million in 2003 to 530 in 2030 with three in four living in developing countries

Clear my choice





Time left 0:28:27

Improved sanitation, nutrition, and medicine during the Industrial Revolution reduced the spread of infectious diseases is related to this Stage of Epidemiologic Transition

- a. Stage III: Degenerative Diseases
- b. Stage I: Pestilence and Famine
- c. Stage IV: Delayed Degenerative
- d. Stage II: Receding Pandemics

Clear my choice

Next page



A graphic description of the relationship between agent, host, and environment is called ..

- a. Epidemiologic homeostasis.
- b. Epidemiologic imbalance.
- c. Epidemiologic transition.
- d. Epidemiologic triad.

[Clear my choice](#)

5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46

Time left 0

A man with a standard life expectancy of 75 years died at age of 60. When he was at age of 40 had amputation from diabetic foot therefore he spent the rest of his life in a disability state with weight factor of 0.5. The DALYs for this person is:

- a. 20 years
- b. 10 years
- c. 15 years
- d. 25 years

[Clear my choice](#)

[Next page](#)

here to search



A group of 30 adventurist went camping in the black forest, Germany, for 3 weeks. After 1 week, 5 persons developed fever, headache, fatigue and skin rash (erythema migrans) 2 of these 5 person also has joint pain. After 2 weeks, 10 more people had the same symptoms and complain and out of 5 had joint pain too. After 3 weeks, 5 more people had the same symptoms and complain and out of 3 had joint pain. Epidemiologist conducted a detailed interview about what may cause these symptoms, the following factors were reported.

Possible causes	Attack rate
Eating wild mushroom	2%
Eating wild berries	75%
Swimming in the lake	15%
Blood sucking Tick Bite	80%
Using water lake for cooking	60%
Wearing long sleeves shirts and long pants	5%
Applying mosquito/insect repellent chemicals	0%

Disease

incubation period ..

- a. Less than 1 week.
- b. Between 2-3 weeks.
- c. Between 1-2 weeks.
- d. More than 3 weeks.

Which of the followings is not among the activities of descriptive epidemiology:

- a. Generate hypotheses
- b. Assess who is affected by the health problem
- c. Study the course of disease within a geographical region
- d. Test the hypothesis to explain the cause of the epidemic
- e. To plan, provide, and evaluate health services

[Clear my choice](#)

5	6	7	8	9	10	11	12	13	14	15	16	17	18	19
30	31	32	33	34	35	36	37	38	39	40	41	42	43	44

طب مجتمع (1)

My courses / طب مجتمع (1) ش1 / General / Med 410 Midterm

What will happen for an epidemic with reproductive rate of 1.5 ..

- a. Infection will last for long time.
- b. Infection will widely spread.
- c. Infection will not widely spread.

Clear my choice

TOSHIBA



Time left 0:29:11

Question 29

Not yet answered

Marked out of 1.0

A comprehensive tobacco control program should only encourage smokers to quit and help them do it

- a. False
- b. True

[Clear my choice](#)

[Next page](#)

طب مجتمع (1)

My courses / طب مجتمع (1) / General / Med 410 Midterm

19

out of

Most NCD deaths around the world according to WHO is accounted for:

- a. Obesity
- b. Cardiovascular diseases
- c. Hypertension
- d. Diabetes
- e. Smoking

[Clear my choice](#)



TOSHIBA



Which statement is incorrect about surveillance?

- a. Phone calls are example of Active surveillance
- b. Active surveillance provides minimal data on risk factors
- c. Patient discharge records are example of passive surveillance
- d. Because passive surveillance depends on people in different institutions to provide data quality and timeliness are difficult to control
- e. Active surveillance is more expensive than passive

[Clear my choice](#)

Search here to search



18

out of

Time left 0:28:38

About QALYs & DALYs, which statement is incorrect?

- a. They both are measurements to calculate the quality and quantity of life either of individuals or the general population
- b. QALY is considered as a modified version of DALY
- c. Both evaluations have different coverage and interpretations
- d. QALYs is regarded as measuring the 'positive', the quality of life, with the aim to maximize the quality of life; while DALYs are measuring the 'negative' impact of a disease or the years lived in poor health
- e. QALYs and DALYs are tools, providing a single measure of mortality and morbidity, used internationally for assessing health care interventions and treatments

Clear my choice

Quiz

1

10

19

28

37

46

Finish after

Improved sanitation, nutrition, and medicine during the Industrial Revolution reduced the spread of diseases is related to this Stage of Epidemiologic Transition

- a. Stage II: Receding Pandemics
- b. Stage I: Pestilence and Famine
- c. Stage IV: Delayed Degenerative
- d. Stage III: Degenerative Diseases

Clear my choice

gation

3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23
28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48

pt ...

طب مجتمع (1)

rd / My courses / طب مجتمع (1) ش 1 / General / Med 410 Midterm

Time left

5

red.
ed out of

A propagated epidemic curve is usually the result of what type of exposure?

- a. Person to person transmission.
- b. Continuous common source.
- c. Intermittent common source.
- d. Point source.

Clear my choice

Ne

Type here to search



Time left 0:31:24

About the impact of non-communicable diseases on global burden of diseases, which statement is incorrect?

- a. The number of patients with diabetes in the world is expected to increase from 194 Million in 2003 to 530 in 2030 with three in four living in developing countries
- b. By 2020, it is predicted that these diseases will be causing seven out of every 10 deaths in developing countries
- c. In some countries of the Middle East, one in four deaths in adults aged between 35 and 64 years is attributable to diabetes
- d. In developed countries most people with diabetes are above the age of retirement, whereas in developing countries those most frequently affected are aged between 35 and 64

[Clear my choice](#)

[Next page](#)

5
A propagated epidemic curve is usually the result of what type of exposure?

- a. Person to person transmission.
- b. Continuous common source.
- c. Intermittent common source.
- d. Point source.

Clear my choice

Time left 0:32:37

Question **27**

Not yet answered

Marked out of 1.0

Offering follow-up for smokers is related to this step of promoting smoking cessation:

- a. Ask
- b. Assist
- c. Advise
- d. Assess
- e. Arrange

[Clear my choice](#)

[Next page](#)

Quick navigation

1	2	3	4
5	6	7	8
9	10	11	12
13	14	15	16
17	18	19	20
21	22	23	24
25	26	27	28
29	30	31	32

[Refresh attempt](#)

طب مجتمع (1) ش 1

Dashboard / My courses / طب مجتمع (1) ش 1 / General / Med 410 Midterm

Time left 0:32:17

Question 15

Not yet answered

Marked out of 1.0

In the conflagration disease spread context, the best option is to ...

- a. Neutralize new viral variants via vaccination.
- b. Very stringent preventive measure.
- c. All is correct
- d. Develop a therapy for sever cases.

[Clear my choice](#)

[Next page](#)

Mitigation is directed to..

- a. Investigate the natural reservoir of the agent.
- b. Slow the spread of the disease.
- c. Identify the infectious agent.
- d. Understand the source of infection.

[Clear my choice](#)

Time left 0:36:14

Next

طب مجتمع

My courses / طب مجتمع (1) ش 1 / General / Med 410 Midterm


Time

What will happen for an epidemic with reproductive rate of 1 ..

- a. Infection will last for long time.
- b. Infection will widely spread.
- c. Infection will not widely spread.

[Clear my choice](#)

Type here to search



Time left

The proportion of infected persons who develop clinical disease is ..

- a. Infectivity.
- b. Pathogenicity.
- c. Virulence.
- d. Immunogenicity.

[Clear my choice](#)

Time left 0:07:45

Identified or diagnosed cases of the infectious disease represent which part of the epidemiologic iceberg?

- a. The tip level of the pyramid.
- b. The bottom base of the pyramid.
- c. The top levels of the pyramid.
- d. The middle levels of the pyramid.

[Clear my choice](#)

Question 6

Not yet answered

Marked out of

1.0

Navigation

27	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
28	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	
29	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25		
30	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25			
31	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25				
32	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25					
33	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25						
34	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25							
35	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25								
36	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25									
37	12	13	14	15	16	17	18	19	20	21	22	23	24	25										
38	13	14	15	16	17	18	19	20	21	22	23	24	25											
39	14	15	16	17	18	19	20	21	22	23	24	25												
40	15	16	17	18	19	20	21	22	23	24	25													
41	16	17	18	19	20	21	22	23	24	25														
42	17	18	19	20	21	22	23	24	25															
43	18	19	20	21	22	23	24	25																
44	19	20	21	22	23	24	25																	
45	20	21	22	23	24	25																		
46	21	22	23	24	25																			
47	22	23	24	25																				
48	23	24	25																					
49	24	25																						
50	25																							

[Next page](#)

Time left 0:38:03

Question 20

Not yet answered

Marked out of 1.0

Indicator: A measurable factor that allows decision makers to estimate objectively the size of a health problem and monitor the processes, the products, or the effects of an intervention on the population

Select one:

True

False

Next page

Quiz n

1 2

10 11

19 20

28 29

37 38

46 47


Finish attempt



An epidemic curve with one peak is usually the result of what type of exposure?

- a. Continuous common source.
- b. Point source.
- c. Person to person transmission.
- d. Intermittent common source.

[Clear my choice](#)



Time left 0:41:18

In the study about mental health status of the general population, healthcare professionals, and university students during the COVID-19 outbreak, the prevalence of depression and anxiety among the entire study participants was 23.8% and 13.1%, respectively

Select one:

- True
- False

1) طب مجتمع

/ My courses / طب مجتمع (1) / General / Med 410 Midterm

Time left 0:42

Indicator: A non measurable factor that allows decision makers to estimate objectively the size of a health problem and monitor the processes, the products, or the effects of an intervention on the population

Select one:

- True
- False

Next page



TOSHIBA



Time left 0:40:29

Question **15**
Not yet answered
Marked out of 1.0

To maximize the success of any program to reduce tobacco use, training for health professionals are related to efforts

- a. Community-Based Programs
- b. Public Education Efforts
- c. Monitoring and Evaluation
- d. Enforcement
- e. Related Policy Efforts

[Clear my choice](#)

[Next page](#)

Quiz navigation

1	2	3	4	5
10	11	12	13	14
19	20	21	22	23
28	29	30	31	32
37	38	39	40	41
46	47	48	49	50

[Finish attempt ...](#)

Time left 0:42:11


A group of 30 adventurist went camping in the black forest, Germany, for 3 weeks. persons developed fever, headache, fatigue and skin rash (erythema migrans) 2 of has joint pain. After 2 weeks, 10 more people had the same symptoms and compl joint pain too. After 3 weeks, 5 more people had the same symptoms and compl joint pain. Epidemiologist conducted a detailed interview about what may caus following factors were reported.

Possible causes	Attack rate
Eating wild mushroom	2%
Eating wild berries	75%
Swimming in the lake	15%
Blood sucking Tick Bite	80%
Using water lake for cooking	60%
Wearing long sleeves shirts and long pants	5%
Applying mosquito/insect repellent chemicals	0%

Source of infection ..

- a. Allergic reaction due to eating berries.
- b. Contaminated water.
- c. Mosquito- borne disease.
- d. Tick borne disease.

Clear my choice



Time left 0:40:41

Benefits of Smoking Cessation include the following except:

- a. Within two weeks to three months of quitting, circulation improves and lung function increases by up to 30 percent
- b. The risk of a major coronary event reduces to the level of a never smoker after 10-15 years
- c. Within two days of quitting, smell and taste are enhanced
- d. Excess risk of heart disease is reduced by half after one year's abstinence
- e. Within one day of quitting, the chance of a heart attack decreases

[Clear my choice](#)

Quiz na

1 2

10 11

19 20

28 29

37 38

46 47

Finish attempt



Time left 0:41:42

In the study about mental health status of the general population, healthcare professionals, and university students during the COVID-19 outbreak, the prevalence of depression and anxiety among the entire study participants was 23.8% and 13.1%, respectively

Select one:

- True
 False

Next page

Quiz n

1	
10	
19	
28	
37	
46	



Finis

Prevalence is the actual number of cases alive and dead

Select one:

True

False



طب مجتمع (1) ش 1

Dashboard / My courses / طب مجتمع (1) ش 1 / General / Med 410 Midterm

Question 7
Not yet answered
Marked out of 1.0

Time left 0:43:35

Develop interventions to reduce the long-term negative impact of low childhood socioeconomic status on cognitive ability and mental health is a challenge that is related related to this goal:

- a. Improve treatments and expand access to care
- b. Raise awareness of the global burden of mental health disorders
- c. Advance prevention and implementation of early interventions
- d. Identify root causes, risk, and protective factors

[Clear my choice](#)

[Next page](#)

Quiz navigation

1	2	3	4	5	6	7	8	9
10	11	12	13	14	15	16	17	18
19	20	21	22	23	24	25	26	27
28	29	30	31	32	33	34	35	36
37	38	39	40	41	42	43	44	45
46	47	48	49	50				

[Finish attempt...](#)

Benefits of Smoking Cessation include the following except:

- a. Within two weeks of quitting, smell and taste are enhanced
- b. Excess risk of heart disease is reduced by half after one year's abstinence
- c. Within one day of quitting, the chance of a heart attack decreases
- d. Within two weeks to three months of quitting, circulation improves and lung function increases by up to 30 percent
- e. The risk of a major coronary event reduces to the level of a never smoker within five years

Clear my choice


A group of 30 adventurist went camping in the black forest, Germany, for 3 weeks. After 1 week, 5 persons developed fever, headache, fatigue and skin rash (erythema migrans) 2 of these 5 person also has joint pain. After 2 weeks, 10 more people had the same symptoms and complain and out of 5 had joint pain too. After 3 weeks, 5 more people had the same symptoms and complain and out of 3 had joint pain. Epidemiologist conducted a detailed interview about what may cause these symptoms, the following factors were reported.]


Possible causes	Attack rate
Eating wild mushroom	2%
Eating wild berries	75%
Swimming in the lake	15%
Blood sucking Tick Bite	80%
Using water lake for cooking	60%
Wearing long sleeves shirts and long pants	5%
Applying mosquito/insect repellent chemicals	0%

Possible

Mode of transmission .

- a. Vector-Borne
- b. Food-borne.
- c. Direct contact with infected individual.
- d. Water-borne.

[Clear my choice](#)



1) طب مجتمع

/ My course / طب مجتمع (1) ش.1 / General / 410 Midterm

Time

The equilibrium between agent, host, and environmental factors is called ...

- a. Epidemiologic homeostasis.
- b. Epidemiologic triad.
- c. Epidemiologic triangle.
- d. Epidemiologic transition.

Clear my choice

TOSHIBA



Which statement is incorrect about Fagerström Test for Nicotine Dependence?

- a. A smoker who smokes 20 cigarettes /day will get 1 point on the scale
- b. Score ranges between 0-10
- c. A score of 7 denotes moderate dependence (addiction)
- d. Contains 5 questions

[Clear my choice](#)

[Next page](#)

Question 5

Not yet answered

Marked out of 1.0

Time left 0:46:01

According to CDC, functions of epidemiology include the followings except

- a. Outbreak Investigation
- b. Evaluation of disease control programs
- c. All options are correct
- d. Offer Interventions
- e. Public Health Surveillance

[Clear my choice](#)

[Next page](#)


Quiz navigation

1	2	3	4
10	11	12	13
19	20	21	22
28	29	30	31
37	38	39	40
46	47	48	49

Finish attempt ...

You are logged in as [ماريا ايمن عيسى المفاريه](#) (Log out)
MED 410 1

Based on your understanding of the natural history of disease spectrum, point (D) represent ..



- a. Onset of symptoms.
- b. Diagnosis time.
- c. Pathologic changes.
- d. Exposure.

[Clear my choice](#)

طب مجتمع (1) ش 1

Dashboard / My courses / طب مجتمع (1) ش 1 / General / Med 410 Midterm

Time left 0:47:16

Question 4
Not yet
answered
Marked out of
1.0

Multiple paid media (TV, radio, print, etc.), public relations, special events and promotions are related to efforts to maximize the success of any program to reduce tobacco use

- a. Monitoring and Evaluation
- b. Public Education Efforts
- c. Related Policy Efforts
- d. Community-Based Programs


[Clear my choice](#)

[Next page](#)

Quiz navigation

1	2	3	4	5
10	11	12	13	14
19	20	21	22	23
28	29	30	31	32
37	38	39	40	41
46	47	48	49	50

Finish attempt



طب مجتمع (1) ش 1

Dashboard / My courses / طب مجتمع (1) ش 1 / General / Med 410 Midterm

Time left 0:48:11

Quiz navigation

1	2	3	4	5
10	11	12	13	14
19	20	21	22	23
28	29	30	31	32
37	38	39	40	41
46	47	48	49	50

Finish attempt...

Next page

One of the indirect way of agent transmission ..

- a. Prenatal from mother to child.
- b. Touching.
- c. Airborne.
- d. Droplets for short distance.

Clear my choice

Question 3

Not yet answered

Marked out of 1.0


Time left 0:49:00

Question 5

Not yet answered

Marked out of 1.0

Based on your understanding of the natural history of disease spectrum, point (C) represent ..



- a. Pathologic changes.
- b. Onset of symptoms.
- c. Diagnosis time.
- d. Exposure.

[Clear my choice](#)

[Next page](#)

Quiz navigation

1	2	3	4	5
10	11	12	13	14
19	20	21	22	23
28	29	30	31	32
37	38	39	40	41
46	47	48	49	50

[Finish attempt ...](#)

Time left

Which of the following infectious diseases may have the largest incidence rate?


- a. Flu.
- b. Common cold.
- c. HIV.
- d. COVID. 19.

Clear my choice

Next

Time l

Based on your understanding of the natural history of disease spectrum, point (B) represent ..



- a. Diagnosis time.
- b. Pathologic changes.
- c. Onset of symptoms.
- d. Exposure.

Clear my choice



TOSHIBA



Syndromic surveillance system can be only active

Select one:

True

False

Time left 0:49:34

Question 2

Not yet answered

Marked out of 1.0

About mental health, which statement is incorrect?

- a. Mental illness occurs in 1 out of every 4 families in the U.S
- b. About 25% of most communities have mental illnesses
- c. -Mental illness can cause chronic diseases
- d. Research output in the Arab world has increased by almost 60% in the past ten years
- e. Mental health includes our emotional, psychological, and social well-being

[Clear my choice](#)

[Next page](#)

Quiz navig

1	2	3
10	11	12
19	20	21
28	29	30
37	38	39
46	47	48

Finish attempt

Time left 0:52:53

Benefits of Smoking Cessation include the following except:

- a. Within two weeks to three months of quitting, circulation improves and lung function increases by up to 30 percent
- b. Excess risk of heart disease is reduced by half after one year's abstinence
- c. The risk of a major coronary event reduces to the level of a never smoker after 10-15 years
- d. Within one day of quitting, the chance of a heart attack decreases
- e. Within two days of quitting, smell and taste are enhanced

[Clear my choice](#)

[Next page](#)

Time left 0:52:04

Question 2
Not yet answered
Marked out of 1.0

Evaluation of surveillance & response systems serves to achieve the followings Except:

- a. Support functions of the system
- b. Document the status of, and any change in the performance of the system
- c. Provide an evidence-base on which to modify surveillance objectives, implementation strategy and planned activities
- d. Provide explanations for achievements and failures in the system
- e. Enable planning of resource allocation

[Clear my choice](#)

Quiz navigation

1	2	3	4	5	6
10	11	12	13	14	15
19	20	21	22	23	24
28	29	30	31	32	33
37	38	39	40	41	42
46	47	48	49	50	

[Finish attempt](#)

[Next page](#)

1) طب مجتمع

/ My courses / طب مجتمع (1) ش1 / General / Med 410 Midterm

An example of vector-borne agent transmission is..

- a. Touching.
- b. Droplets for short distance.
- c. Mosquito-borne.
- d. Prenatal from mother to child.

[Clear my choice](#)



TOSHIBA



Time left 0:54:14

Which of the following intervals or time period determine when person is capable of infecting other people?

- a. Infectious period.
- b. Latent period.
- c. Incubation period.
- d. Serial interval.

[Clear my choice](#)

[Next page](#)

- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26

Time left 0:52:20

Question 1
Not yet answered
Marked out of 1.0

About the impact of non-communicable diseases on global burden of diseases, which statement is incorrect?

- a. In some countries of the Middle East, one in four deaths in adults aged between 35 and 64 years is attributable to diabetes
- b. In developed countries most people with diabetes are above the age of retirement, whereas in developing countries those most frequently affected are aged between 35 and 64
- c. By 2020, it is predicted that these diseases will be causing seven out of every 10 deaths in developing countries
- d. The number of patients with diabetes in the world is expected to increase from 194 Million in 2003 to 530 in 2030 with three in four living in developing countries

[Clear my choice](#)

[Next page](#)

Quiz navigation

1	2	3	4	5	6	7	8	9
10	11	12	13	14	15	16	17	18
19	20	21	22	23	24	25	26	27
28	29	30	31	32	33	34	35	36
37	38	39	40	41	42	43	44	45
46	47	48	49	50				

[Finish attempt ...](#)

Question 1

Not yet answered
Marked out of

Time left 0:53:18

About QALYs & DALYs, which statement is incorrect?

- a. QALYs is regarded as measuring the 'positive', the quality of life, with the aim to maximize the quality of life; while DALYs are measuring the 'negative' impact of a disease or the years lived in poor health
- b. DALY is considered as a modified version of QALY
- c. They both are measurements to calculate the quality and quantity of life either of individuals or the general population
- d. QALYs and DALYs are tools, providing a single measure of mortality and morbidity, used internationally for assessing health care interventions and treatments
- e. Both evaluations have similar coverage and interpretations

[Clear my choice](#)

1	2
10	11
19	20
28	29
37	38
46	47

[Finish attempt](#)

Time left 0:53:52

About WHO STEPwise approach to Risk Factor Surveillance (STEPS), which statement is incorrect?

- a. The WHO STEPwise approach to Surveillance (STEPS) is a simple, standardized method for collecting, analysing and disseminating data in WHO member countries
- b. Since its inception, the STEPS approach has advocated that large amounts of poor-quality data are more valuable than small amounts of good-quality data
- c. The WHO STEPwise approach to noncommunicable disease (NCD) risk factor surveillance is designed to help countries build and strengthen their surveillance capacity
- d. The STEPS approach supports monitoring a few modifiable NCD risk factors that reflect a large part of the future NCD burden
- e. The approach encourages the collection of small amounts of useful information on a regular and continuing basis

[Clear my choice](#)

Quiz na

1	2
10	11
19	20
28	29
37	38
46	47

Finish atten

