## Coronavirus Disease 2019 (COVID-19) Personal Protective Equipment (PPE) Guidelines

Applicable to PPE use in the hospital setting, including the emergency department, and ambulatory settings

## Current as of November 17, 2020

**Northwell** 

Health<sup>®</sup>

These guidelines apply in cases when PPE is worn for individual patient interactions.

PPE	Use	Details
Gowns	1	Impervious/Fluid-resistant: Blood or body fluid risk, including aerosol-generating procedures (AGPs) such as intubation, regardless of patient's COVID-19 status. Remove if contaminated.Isolation: When there is no blood or body fluid risk. Remove if contaminated.
Eye protection (mask with face shield, face shield or goggles)	1	Use with suspected or confirmed COVID-19 patients, for all AGPs, and for all encounters with any individual who is unable to wear a mask. If reusing, disinfect with an Environmental Protection Agency (EPA) approved wipe, air dry, and place in labeled bag with name.
Procedure mask	<b>√</b>	Required to be worn at all times in the hospital, ambulatory or practice office settings.Place over N95 respirator: - When caring for a suspected or confirmed COVID-19 case without aerosols.Not required over N95 respirator for AGPs for suspected or confirmed COVID-19 patients since N95 respirator will be discarded.Patients should wear a procedure mask when outside their room and in their room whenever possible.Place over N95 respirator: - When caring for a suspected or confirmed COVID-19 case 
N95 respirator	1	Worn for all patient encounters where individuals have suspected or confirmed COVID-19 status. Worn for all AGPs regardless of patient's COVID-19 status. Discard ONLY after suspected or confirmed COVID-19 patient contact. Between uses, store in breathable bag labeled with name.