Coronavirus Disease 2019 (COVID-19)



Personal Protective Equipment (PPE) Reuse & Replace Guidelines

Current as of March 3, 2021

Applicable to PPE use in the hospital setting, including the emergency department, and ambulatory setting These guidelines apply in cases when PPE is worn for individual patient interactions.

Upon request by those interacting directly with patients, we will replace uncontaminated N95 respirators once each day.

Reuse Replace		Suspected or confirmed COVID-19 case WITHOUT aerosols	For all patients WITH aerosol-generating procedures
	N95 respirator	Reuse Place in a breathable bag labeled with your name*	Replace after completion of the aerosol-generating procedure
	Procedure mask	N/A	N/A
Ċ	External cloth mask over procedure mask	N/A	N/A
	Procedure mask with attached face shield	N/A	N/A
	Eye protection	Reuse Disinfect with an Environmental Protection Agency (EPA) approved wipe. Then air dry and place in a bag labeled with your name.	Reuse Disinfect with an Environmental Protection Agency (EPA) approved wipe. Then air dry and place in a bag labeled with your name.
	Impervious/ Fluid-resistant gown	Replace after each patient encounter involving blood or body fluid	Replace after completion of the aerosol-generating procedure
	Isolation gown	Replace after each patient encounter	N/A
JUM	Gloves	Replace	Replace

NOTE: Replace any PPE that becomes contaminated *Reuse Protocol: See "Recommended Guidance for Extended Use and Limited Re-Use of N95 Filtering Facepiece Respirators in Healthcare Settings" on the Emergency Management Novel Coronavirus (COVID-19) Information webpage.